

Print selected from 10562215.trn

=> s ll full
FULL SEARCH INITIATED 11:16:52 FILE 'CASREACT'
SCREENING COMPLETE - 6095 REACTIONS TO VERIFY FROM 502 DOCUMENTS

100.0% DONE 6095 VERIFIED 1658 HIT RUNS (120 INCOMP) 114 DOCS
SEARCH TIME: 06:00.04

L3 114 SEA SSS FUL L1 (1458 REACTIONS)

=> s l3 and carbon monoxide
50134 CARBON
1717 CARBONS
51225 CARBON
(CARBON OR CARBONS)
6370 MONOXIDE
92 MONOXIDES
6411 MONOXIDE
(MONOXIDE OR MONOXIDES)
5849 CARBON MONOXIDE
(CARBON(W)MONOXIDE)
L4 0 L3 AND CARBON MONOXIDE

=> s carbon monoxide
50134 CARBON
1717 CARBONS
51225 CARBON
(CARBON OR CARBONS)
6370 MONOXIDE
92 MONOXIDES
6411 MONOXIDE
(MONOXIDE OR MONOXIDES)
L5 5849 CARBON MONOXIDE
(CARBON(W)MONOXIDE)

=> file reg
FILE 'REGISTRY' ENTERED AT 11:17:25 ON 16 MAR 2009
USE IS SUBJECT TO THE TERMS OF YOUR SIN CUSTOMER AGREEMENT.
PLEASE SEE 'HELP USAGETERMS' FOR DETAILS.
COPYRIGHT (C) 2009 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 13 MAR 2009 HIGHEST RN 1120564-02-4
DICTIONARY FILE UPDATES: 13 MAR 2009 HIGHEST RN 1120564-02-4

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 9, 2009.

Please note that search-term pricing does apply when
conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and
predicted properties as well as tags indicating availability of
experimental property data in the original document. For information
on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/scndoc/properties.html>

Print selected from 10562215.trn

=> s carbon monoxide/cn
L6 1 CARBON MONOXIDE/CN

=> d rcd

L6 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2009 ACS ON SIN

=> d rn

L6 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2009 ACS ON SIN
RN 630-08-0 REGISTRY

=> file casreact
FILE 'CASREACT' ENTERED AT 11:17:52 ON 16 MAR 2009
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is
held by the publishers listed in the PUBLISHER (PB) field (available
for records published or updated in Chemical Abstracts after December
26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 15 MAR 2009 VOL 150 ISS 12

New CAS Information Use Policies, enter HELP USAGETERMS for details.

* CASREACT now has more than 16.5 million reactions *
* *****

CASREACT contains reactions from CAS and from: ZIC/VINITI database
(1974-1999) provided by InfoChem; INPI data prior to 1986;
Biotransformations database compiled under the direction of
Professor Dr. Klaus Kieselich; organic reactions, portions copyright
1996-2006 John Wiley & Sons, Ltd., John Wiley and Sons, Inc.,
Organic Reactions Inc., and Organic Syntheses Inc. Reproduced under
license. All Rights Reserved.

This file contains CAS Registry Numbers for easy and accurate substance
identification.

=> s 630-08-0
L7 10878 630-08-0

=> d his

(FILE 'HOME' ENTERED AT 11:15:49 ON 16 MAR 2009)

FILE 'CASREACT' ENTERED AT 11:16:04 ON 16 MAR 2009
L1 STRUCTURE UPLOADED
L2 3 S L1
L3 114 S L1 FULL
L4 0 S L3 AND CARBON MONOXIDE
L5 5849 S CARBON MONOXIDE

Print selected from 10562215.trn

FILE 'REGISTRY' ENTERED AT 11:17:25 ON 16 MAR 2009
1 S CARBON MONOXIDE/CN




FILE 'CASREACT' ENTERED AT 11:17:52 ON 16 MAR 2009
10878 S 630-08-0

=> s 17 and 13
L8 2 L7 AND L3

=> d chib abs ord 1-
YOU HAVE REQUESTED DATA FROM 2 ANSWERS - CONTINUE? Y(N):y

18 ANSWER 1 OF 2 CASREACT COPYRIGHT 2009 ACS on STN
144:292572 Diaminocoumarins as fluorogenic substrates for monoamine oxidases,
and their preparation, pharmaceutical compositions, photophysical
properties, and a method for detecting active monoamine oxidases and their
inhibitors for treatment of nervous system disorders. Chen, Song; Yee,
Dominic J.; Subramanian, Mike; Sames, Dalibor [The Trustees of Columbia
University in the City of New York, USA]. PCT Int. Appl. NO 2006/026368
A2 20060309, 278 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BW, BY, BE, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ,
EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HU, HR, ID, IL, IN, IS, JP, KE,
KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, NZ, NA, NG, NI, NO, NZ, OM, EG, PH, PL, PT, RO, RU, SC, SE, SG, SK,
SL, SR, SY, TJ, TM, TN, TR, TT, TZ, UA, US, UZ, VC, VN, YU, ZA,
ZM, ZW; RW: AT, BE, BF, BJ, CE, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR,
GA, GB, GR, HE, IS, IT, LU, MC, ML, NE, NL, PT, SE, SN, TD, TS, TR.
(English). CODEN: PIXXDC. APPLICATION: NO 2005-US0276 20050825.
PRIORITY: US 2004-604538P 20040825.

GI




AB The invention relates to compds. of formula I, which are useful for
detecting the activity of monoamine oxidases (MAO), compds. useful for
competitively inhibiting monoamine oxidases, for determining inhibitors of
monoamine oxidases and compds. useful for treating monoamine

Print selected from 10562215.trn


oxidase-related nervous system pathologies, as well as pharmaceutical
compos. and methods of manufacture thereof. Compds. of formula I wherein R1 is
H, alkyl, alkenyl, alkynyl, (un)substituted (hetero)aryl, cycloalkyl, NH2
and derivs., alkyl-CO2H, alkyl-CN, alkyl-NH2, halo, CN, or indole; R2-R6
are independently H, OH, alkyl, alkenyl, alkynyl, (un)substituted
(hetero)aryl, cycloalkyl, NH2 and derivs., alkyl-CO2H, alkyl-CN,
alkyl-NH2, O-alkyl, O-alkenyl, O-alkynyl, O-aryl, O-cycloalkyl, CN, halo,
or indole; R1R2 and R1R6 may independently form an unsubstituted pyrrole;
R1R2R6 may form an octahydroquinazolinone; R2R3 may form a pyrrole; X is
halo; or pharmaceutically acceptable salts, or stereoisomers thereof are
claimed in this invention. The process for preparing these fluorogenic
substrates, and a method for identifying a test compound as a substrate of
MAO are also claimed. Example compound II was prepared by nitration of
7-methylcoumarin, and the resulting 7-methyl-6-nitrocoumarin underwent
condensation with DMF di-Na acetal to give the corresponding
coumarin-enamine, which was hydrolyzed to give
7-(2-methyl-1-6-nitrocoumarin, which underwent reductive amination with
dibenzosuberilamine; the resulting
7-(dibenzosuberilaminomethyl)-6-nitrocoumarin was reduced to the
6-amminocoumarin compound, which was hydrolyzed to give coumarin II. The
invention compds. were evaluated for their photophys. properties and their
enzymic activity toward MAO-A and MAO-B. Coumarin II showed λ_{max}
352 nm, ϵ 2000±100 M⁻¹cm⁻¹, ϕ 0.001±0.0001, and
k_{em} was not detected. The invention compds. were evaluated for
their ability to convert to the corresponding indole in the presence of
MAO-A and MAO-B. Enzyme-catalyzed indole formation was realized after 24
h by reading fluorescence emission upon excitation at the absorbance
maxima of the resp. indoles. Only diaminocoumarin II showed a
significant conversion to its indole III in the presence of MAO-A and
MAO-B. The enzyme kinetics for fluorogenic probe II were also determined
Compound II showed for MAO-A: K_m = 30.9±1.8 μ M, K_{cat} = 0.475 min⁻¹,
K_{cat}/K_m = 0.015 min⁻¹ μ M⁻¹, V_{max}(mitochondria) = 0.0570 nmol min⁻¹
mg⁻¹; and for MAO-B: K_m = 510±40 μ M, K_{cat} = 20.62 min⁻¹, K_{cat}/K_m =
0.040 min⁻¹ μ M⁻¹, and V_{max}(mitochondria) = 1.2 nmol min⁻¹ mg⁻¹. These
results indicate that coumarin probe II is a good substrate for monoamine
oxidase and could be useful as a fluorescent reporter for monoamine
oxidase.

RX(15) OF 512



CON: 2 hours, room temperature

RX(21) OF 512





(step 1)

CON: STAGE(1) 50 deg C
STAGE(2) 2 hours, 50 deg C


Print selected from 10562215.trn

RX(107) OF 512 - 2 STEPS




NOTE: 2) fuming nitric acid used
 CON: STEP(1.1) 50 deg C
 STEP(1.2) 2 hours, 50 deg C
 STEP(2) 2 hours, 0 deg C

RX(181) OF 512 - 2 STEPS



CON: STEP(1.1) room temperature; 12 hours, reflux
 STEP(2.1) 50 deg C
 STEP(2.2) 2 hours, 50 deg C


RX(201) OF 512 - 3 STEPS



NOTE: 2) fuming nitric acid used
 CON: STEP(1.1) 50 deg C
 STEP(1.2) 2 hours, 50 deg C
 STEP(2) 2 hours, 0 deg C
 STEP(3) 1 hour, 110 deg C



Print selected from 10562215.trn

RX(202) OF 512 - 4 STEPS



NOTE: 3) fuming nitric acid used
 CON: STEP(1.1) room temperature; 12 hours, reflux
 STEP(2.1) 50 deg C
 STEP(2.2) 2 hours, 50 deg C
 STEP(3) 2 hours, 0 deg C
 STEP(4) 1 hour, 110 deg C


RX(317) OF 512 - 3 STEPS




NOTE: 3) fuming nitric acid used
 CON: STEP(1.1) room temperature; 12 hours, reflux
 STEP(2.1) 50 deg C
 STEP(2.2) 2 hours, 50 deg C
 STEP(3) 2 hours, 0 deg C

Print selected from 10562215.trn

RX(388) OF 512 - 8 STEPS



RX(388) OF 512 - 8 STEPS




NOTE: 6) Suzuki coupling in stage 2, key step in stage 2, 8)
Dess-Martin oxidation

CON: STEP(1.1) 0 deg C; 1 hour, 0 deg C
STEP(2.1) 0 deg C; overnight, room temperature
STEP(3) 4 hours, room temperature
STEP(4) 20 hours, reflux
STEP(5) 4 hours, 100 deg C
STEP(6.1) 2 hours, room temperature
STEP(6.2) 20 hours, 80 deg C
STEP(7.1) room temperature; 30 minutes, room temperature
STEP(8.1) room temperature; 30 minutes, room temperature

Print selected from 10562215.trn

RX(497) OF 512 - 9 STEPS



NOTE: 6) Suzuki coupling in stage 2, key step in stage 2, 8)
Dess-Martin oxidation

CON: STEP(1.1) 0 deg C; 1 hour, 0 deg C
STEP(2.1) 0 deg C; overnight, room temperature
STEP(3) 4 hours, room temperature
STEP(4) 20 hours, reflux
STEP(5) 4 hours, 100 deg C
STEP(6.1) 2 hours, room temperature
STEP(6.2) 20 hours, 80 deg C
STEP(7.1) room temperature; 30 minutes, room temperature
STEP(8.1) room temperature; 30 minutes, room temperature
STEP(9) 30 minutes, room temperature

LB ANSWER 2 OF 2 CASREACT COPYRIGHT 2009 ACS on STN

142:411180 Synthesis of 5-Substituted-1H-indol-2-yl-1H-quinolin-2-ones: A Novel Class of KDR Kinase Inhibitors. Kuehe, Jeffrey T.; Wong, Audrey; Qi, Chuanming; Smitrovich, Jacqueline; Davies, Ian W.; Hughes, David L. (Department of Process Research, Merck & Co., Inc., Rahway, NJ, 07065, USA). Journal of Organic Chemistry, 70(7), 2555-2567 (English) 2005. CODEN: JOCEAH. ISSN: 0022-3263. Publisher: American Chemical Society.

GI


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB A number of approaches for the synthesis of the 1H-indol-2-yl-1H-quinolin-2-one ring system found in the potent and selective KDR kinase inhibitor I are described. The preparation and reaction of trimethylsilylnitrobenzene II with 2-methoxy-3-quinolinecarboxaldehyde afforded alc. III, which was the key intermediate for the preparation of the target compds. Conversion of alc. III to either nitroketone IV or


Print selected from 10562215.trn

nitrostyrene V set the stage for reductive cyclization. The quinolin-2-one functionality was unmasked in the last step to provide compound I in 56-60% overall yield from readily available starting materials.

RX(22) OF 350




RX(22) OF 350



NOTE: Raney Nickel used
CON: 7.5 hours, 65 deg C, 40 psi


Print selected from 10562215.trn

RX(27) OF 350




NOTE: Raney Nickel used
CON: 7.5 hours, 65 deg C, 40 psi

RX(28) OF 350





RX(28) OF 350



NOTE: Raney Nickel used
CON: 20 hours, room temperature, 40 psi

Print selected from 10562215.trn




RX(36) CF 350



94%

CON: 14 hours, 70 deg C, 15 psi


RX(37) CF 350



95%



Print selected from 10562215.trn

RX(37) CF 350



CON: 15 hours, 70 deg C, 40 atm

RX(58) CF 350 - 2 STEPS




HCl
100%

NOTE: 1) Raney Nickel used

CON: STEP(1) 7.5 hours, 65 deg C, 40 psi


Print selected from 10562215.trn

RX(59) OF 350 - 2 STEPS



1. Pd(OAc)₂,
1,10-Phenanthroline,
CO, DMF
2. HCl, Water, DMF


RX(59) OF 350 - 2 STEPS



HCl
100%

CON: STEP(1) 14 hours, 70 deg C, 15 psi


RX(60) OF 350 - 2 STEPS



1. Pd(OAc)₂, PPh₃, CO,
MeCN
2. HCl, Water, DMF

Print selected from 10562215.trn



RX(60) OF 350 - 2 STEPS



HCl
100%

CON: STEP(1) 15 hours, 70 deg C, 60 atm

RX(79) OF 350 - 2 STEPS



1. Ni, H₂, EtOH, THF
2. AcOH, Water


HCl
93%

NOTE: 1) Raney Nickel used

CON: STEP(1) 20 hours, room temperature, 40 psi
STEP(2) reflux


Print selected from 10562215.trn

RX(81) OF 350 - 2 STEPS



1.1. (CF₃CO)₂O,
S:108-21-4
1.2. DBU
2. Pd(OAc)₂,
1,10-Phenanthroline,
CO, DMF

RX(81) OF 350 - 2 STEPS




94%

NOTE: 1) stereoselective

CON: STEP(1.1) 30 minutes, room temperature
STEP(1.2) 30 minutes, 60 deg C
STEP(2) 14 hours, 70 deg C, 15 psi


RX(82) OF 350 - 2 STEPS




1.1. (CF₃CO)₂O,
S:108-21-4
1.2. DBU
2. Pd(OAc)₂, PPh₃, CO,
MeCN

Print selected from 10562215.trn

RX(82) OF 350 - 2 STEPS




95%



NOTE: 1) stereoselective

CON: STEP(1.1) 30 minutes, room temperature
STEP(1.2) 30 minutes, 60 deg C
STEP(2) 15 hours, 70 deg C, 60 atm


RX(84) OF 350 - 2 STEPS



1. Tri-*o*-tolyphosphine,
Pd(OAc)₂, Et₃N,
DMF
2. Pd(OAc)₂,
1,10-Phenanthroline,
CO, DMF

Print selected from 10562215.trn


RX(84) OF 350 - 2 STEPS



NOTE: 1) stereoselective

CON: STEP(1) 4 hours, 100 deg C
STEP(2) 14 hours, 70 deg C, 15 psi



RX(85) OF 350 - 2 STEPS



1. Tri-*o*-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
2. Pd(OAc)₂, PPh₃, CO,
MeCN

Print selected from 10562215.trn


RX(85) OF 350 - 2 STEPS



NOTE: 1) stereoselective

CON: STEP(1) 4 hours, 100 deg C
STEP(2) 15 hours, 70 deg C, 60 atm


RX(115) OF 350 - 3 STEPS



- 1.1. Na.(AcO) 3BH,
CH₂Cl₂
- 1.2. NH₄Cl, Water
2. Tri-*o*-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
3. Pd(OAc)₂,
1,10-Phenanthroline,
CO, DMF

Print selected from 10562215.trn

RX (115) OF 350 - 3 STEPS



NOTE: 2) stereoselective


CON: STEP (1.1) 4 hours, room temperature

STEP (1.2) room temperature

STEP (2) 4 hours, 100 deg C

STEP (3) 14 hours, 70 deg C, 15 psi

RX (116) OF 350 - 3 STEPS



1.1. Na. (AcO) 3BB,
CHCl₂

1.2. NH₄Cl, Water

2. Tri-*o*-tolylphosphine,

Pd(OAc)₂, Et₃N,


DMF

3. Pd(OAc)₂, PPh₃, CO,

MeCN

Print selected from 10562215.trn

RX (116) OF 350 - 3 STEPS



NOTE: 2) stereoselective


CON: STEP (1.1) 4 hours, room temperature

STEP (1.2) room temperature

STEP (2) 4 hours, 100 deg C

STEP (3) 15 hours, 70 deg C, 60 atm

RX (122) OF 350 - 3 STEPS



1. C: 98327-87-8,

R: 338746-12-6, KBr,

Water, AcOH

2. Tri-*o*-tolylphosphine,

Pd(OAc)₂, Et₃N,

DMF


3. Pd(OAc)₂,

1,10-Phenanthroline,

CO, DMF

Print selected from 10562215.trn


RX(122) OF 350 - 3 STEPS



NOTE: 2) stereoselective

CON: STEP(1) 4 hours, room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 14 hours, 70 deg C, 15 psi



RX(123) OF 350 - 3 STEPS



1. C:98327-87-8,
R:338746-12-6, KBr,
Water, AcOH
2. Tri-o-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
3. Pd(OAc)₂, PPh₃, CO,
MeCN

Print selected from 10562215.trn


RX(123) OF 350 - 3 STEPS



NOTE: 2) stereoselective

CON: STEP(1) 4 hours, room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 15 hours, 70 deg C, 60 atm


RX(129) OF 350 - 4 STEPS



1. Pd, H₂, AcOEt
2. C:98327-87-8,
R:338746-12-6, KBr,
Water, AcOH
3. Tri-o-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
4. Pd(OAc)₂,
1,10-Phenanthroline,
CO, DMF

Print selected from 10562215.trn

RX(129) OF 350 - 4 STEPS




94%

NOTE: 3) stereoselective

CON: STEP(1) 4 hours, room temperature
STEP(2) 4 hours, room temperature
STEP(3) 4 hours, 100 deg C
STEP(4) 14 hours, 70 deg C, 15 psi

RX(130) OF 350 - 4 STEPS




(step 3)


1. Pd, Et, AcOEt
2. C-98327-07-8, KBr,
Water, AcOH
3. Tri-*n*-butylphosphine,
Pd(OAc)₂, Et₃N,
DMF
4. Pd(OAc)₂, PPh₃, CO,
MeCN

Print selected from 10562215.trn

RX(130) OF 350 - 4 STEPS




95%



NOTE: 3) stereoselective

CON: STEP(1) 4 hours, room temperature
STEP(2) 4 hours, room temperature
STEP(3) 4 hours, 100 deg C
STEP(4) 15 hours, 70 deg C, 60 atm


RX(164) OF 350 - 3 STEPS



1. Bu₄N.F, S:108-21-4
- 2.1. (CF₃CO)₂O,
S:108-21-4
- 2.2. DBU
3. Pd(OAc)₂,
1,10-Phenanthroline,
CO, DMF


Print selected from 10562215.trn

RX(164) OF 350 - 3 STEPS



NOTE: 1) key intermediate, 2) stereoselective
 CON: STEP(1.1) room temperature; 30 minutes, room temperature
 STEP(2.1) 30 minutes, room temperature
 STEP(2.2) 30 minutes, 60 deg C
 STEP(3) 14 hours, 70 deg C, 15 psi



RX(165) OF 350 - 3 STEPS



1. Bu4N.F, S:108-21-4
 2.1. (CF3CO)2O,
 S:108-21-4
 2.2. DBU
 3. Pd(OAc)2, PPh3, CO,
 MeCN


Print selected from 10562215.trn

RX(165) OF 350 - 3 STEPS



NOTE: 1) key intermediate, 2) stereoselective
 CON: STEP(1.1) room temperature; 30 minutes, room temperature
 STEP(2.1) 30 minutes, room temperature
 STEP(2.2) 30 minutes, 60 deg C
 STEP(3) 15 hours, 70 deg C, 60 atm


RX(168) OF 350 - 4 STEPS



1.1. Me3SiCH2MgCl,
 Et2O, THF
 1.2. I2, Water
 2. Bu4N.F, S:108-21-4
 3.1. (CF3CO)2O,
 S:108-21-4
 3.2. DBU
 4. Pd(OAc)2,
 1,10-Phenanthroline,
 CO, DMF


Print selected from 10562215.trn

RX(168) OF 350 - 4 STEPS



NOTE: 2) key intermediate, 3) stereoselective
 CON: STEP(1.1) -20 deg C; <-5 deg C; 30 minutes, -20 deg C
 STEP(1.2) 3 hours, room temperature
 STEP(2.1) room temperature; 30 minutes, room temperature
 STEP(3.1) 30 minutes, room temperature
 STEP(3.2) 30 minutes, 60 deg C
 STEP(4) 14 hours, 70 deg C, 15 psi


RX(169) OF 350 - 4 STEPS



1.1. MeSi(CH₃)MgCl,
 Et₂O, THF
 1.2. I₂, Water
 2. Bu₄N.F, S:108-21-4
 3.1. (CF₃CO)₂O,
 S:108-21-4
 3.2. DBU
 4. Pd(OAc)₂, PPh₃, CO,
 MeCN


Print selected from 10562215.trn

RX(169) OF 350 - 4 STEPS



NOTE: 2) key intermediate, 3) stereoselective
 CON: STEP(1.1) -20 deg C; <-5 deg C; 30 minutes, -20 deg C
 STEP(1.2) 3 hours, room temperature
 STEP(2.1) room temperature; 30 minutes, room temperature
 STEP(3.1) 30 minutes, room temperature
 STEP(3.2) 30 minutes, 60 deg C
 STEP(4) 15 hours, 70 deg C, 60 atm


RX(172) OF 350 - 4 STEPS



1. MeOH, KOH
 2. Bu₄N.F, S:108-21-4
 3.1. (CF₃CO)₂O,
 S:108-21-4
 3.2. DBU
 4. Pd(OAc)₂,
 1,10-Phenanthroline,
 CO, DMF

Print selected from 10562215.trn

RX(172) OF 350 - 4 STEPS



NOTE: 2) key intermediate, 3) stereoselective

CON: STEP(1) 2.5 hours, reflux


STEP(2.1) room temperature; 30 minutes, room temperature

STEP(3.1) 30 minutes, room temperature

STEP(3.2) 30 minutes, 60 deg C

STEP(4) 14 hours, 70 deg C, 15 psi

RX(173) OF 350 - 4 STEPS



1. MeOH, KOH

2. Bu4N.F, S:108-21-4


3.1. (CF3CO)2O,
S:108-21-4

3.2. DBU

4. Pd(OAc)2, PPh3, CO,
MeCN

Print selected from 10562215.trn

RX(173) OF 350 - 4 STEPS



NOTE: 2) key intermediate, 3) stereoselective

CON: STEP(1) 2.5 hours, reflux


STEP(2.1) room temperature; 30 minutes, room temperature

STEP(3.1) 30 minutes, room temperature

STEP(3.2) 30 minutes, 60 deg C

STEP(4) 15 hours, 70 deg C, 60 atm

RX(188) OF 350 - 3 STEPS



1.1. (CF3CO)2O,

S:108-21-4

1.2. DBU

2. Pd(OAc)2,


1,10-Phenanthroline,

CO, DMF

3. HCl, Water, DMF

Print selected from 10562215.trn


RX(188) OF 350 - 3 STEPS



HCl
100%

NOTE: 1) stereoselective
CON: STEP(1.1) 30 minutes, room temperature
STEP(1.2) 30 minutes, 60 deg C
STEP(2) 14 hours, 70 deg C, 15 psi


RX(189) OF 350 - 3 STEPS



1.1. (CF₃CO)₂O,
8:108-21-4
1.2. Et₃N
2. Pd(OAc)₂, PPh₃, CO,
MeCN
3. HCl, Water, DMF

Print selected from 10562215.trn


RX(189) OF 350 - 3 STEPS



HCl
100%

NOTE: 1) stereoselective
CON: STEP(1.1) 30 minutes, room temperature
STEP(1.2) 30 minutes, 60 deg C
STEP(2) 15 hours, 70 deg C, 60 atm


RX(190) OF 350 - 3 STEPS



1. Tri-*o*-tolyphosphine,
Pd(OAc)₂, Et₃N,
DMF
2. Pd(OAc)₂,
1,10-Phenanthroline,
CO, DMF
3. HCl, Water, DMF

Print selected from 10562215.trn

RX(190) OF 350 - 3 STEPS




HCl
100%

NOTE: 1) stereoselective


CON: STEP(1) 4 hours, 100 deg C
STEP(2) 14 hours, 70 deg C, 15 psi

RX(191) OF 350 - 3 STEPS



1. Tri-*o*-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
2. Pd(OAc)₂, PPh₃, CO,
MeCN
3. HCl, Water, DMF

RX(191) OF 350 - 3 STEPS




HCl
100%

NOTE: 1) stereoselective

CON: STEP(1) 4 hours, 100 deg C
STEP(2) 15 hours, 70 deg C, 60 atm


Print selected from 10562215.trn

RX(192) OF 350 - 4 STEPS



- 1.1. Na.(AcO)3BH,
CH₂Cl₂
- 1.2. NH₄Cl, Water
2. Tri-*o*-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
3. Pd(OAc)₂,
1,10-Phenanthroline,
CO, DMF
4. HCl, Water, DMF

RX(192) OF 350 - 4 STEPS




HCl
100%

NOTE: 2) stereoselective

CON: STEP(1.1) 4 hours, room temperature
STEP(1.2) room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 14 hours, 70 deg C, 15 psi


Print selected from 10562215.trn

RX(193) CF 350 - 4 STEPS



- 1.1. Na.(AcO)3BH,
CH2Cl2
- 1.2. NH4Cl, Water
2. Tri-o-tolylphosphine,
Pd(OAc)2, Et3N,
DMF
3. Pd(OAc)2, BPh3, CO,
MeCN
4. HCl, Water, DMF

RX(193) CF 350 - 4 STEPS




HCl
100%

NOTE: 2) stereoselective

CON: STEP(1.1) 4 hours, room temperature
STEP(1.2) room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 15 hours, 70 deg C, 60 atm


Print selected from 10562215.trn

RX(194) CF 350 - 4 STEPS



1. C:98327-87-8,
R:338746-12-6, KBr,
Water, AcOH
2. Tri-o-tolylphosphine,
Pd(OAc)2, Et3N,
DMF
3. Pd(OAc)2,
1,10-Phenanthroline,
CC, DMF
4. HCl, Water, DMF

RX(194) CF 350 - 4 STEPS




HCl
100%

NOTE: 2) stereoselective

CON: STEP(1) 4 hours, room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 14 hours, 70 deg C, 15 psi


Print selected from 10562215.trn

RX(195) OF 350 - 4 STEPS



1. C:98327-87-8,
R:338746-12-6, KBr,
Water, AcOH
2. Tri-*o*-tolylphosphine,
 $\text{Pd}(\text{OAc})_2$, Et₃N,
DMF
3. $\text{Pd}(\text{OAc})_2$, PPh₃, CO,
MeCN
4. HCl, Water, DMF

RX(195) OF 350 - 4 STEPS




HCl
100%

NOTE: 2) stereoselective

CON: STEP(1) 4 hours, room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 15 hours, 70 deg C, 60 atm


Print selected from 10562215.trn

RX(196) OF 350 - 4 STEPS



1. Bu₄N.F, S:108-21-4
- 2.1. (CF₃CO)₂O,
S:108-21-4
- 2.2. DBU
3. $\text{Pd}(\text{OAc})_2$,
1,10-Phenanthroline,
CO, DMF
4. HCl, Water, DMF

RX(196) OF 350 - 4 STEPS




HCl
100%

NOTE: 1) key intermediate, 2) stereoselective

CON: STEP(1.1) room temperature; 30 minutes, room temperature
STEP(2.1) 30 minutes, room temperature
STEP(2.2) 30 minutes, 60 deg C
STEP(3) 14 hours, 70 deg C, 15 psi


RX(197) OF 350 - 4 STEPS



1. Bu₄N.F, S:108-21-4
- 2.1. (CF₃CO)₂O,
S:108-21-4
- 2.2. DBU
3. $\text{Pd}(\text{OAc})_2$, PPh₃, CO,
MeCN
4. HCl, Water, DMF

Print selected from 10562215.trn


RX(197) OF 350 - 4 STEPS



HCl
100%

NOTE: 1) key intermediate, 2) stereoselective
CON: STEP(1.1) room temperature, 30 minutes, room temperature
STEP(2.1) 30 minutes, room temperature
STEP(2.2) 30 minutes, 60 deg C
STEP(3) 15 hours, 70 deg C, 60 atm


RX(198) OF 350 - 4 STEPS



- 1.1. BuLi, THF
- 1.2. THF
2. Tri-*o*-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
3. Pd(OAc)₂,
1,10-Phenanthroline,
CO, DMF
4. HCl, Water, DMF

Print selected from 10562215.trn


RX(198) OF 350 - 4 STEPS



HCl
100%

NOTE: 2) stereoselective
CON: STEP(1.1) 30 minutes, room temperature
STEP(1.2) 1 hour, room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 14 hours, 70 deg C, 15 psi


RX(199) OF 350 - 4 STEPS



- 1.1. BuLi, THF
- 1.2. THF
2. Tri-*o*-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
3. Pd(OAc)₂, PPh₃, CO,
MeCN
4. HCl, Water, DMF

Print selected from 10562215.trn

RX(199) OF 350 - 4 STEPS




HCl
100%

NOTE: 2) stereoselective

CON: STEP(1.1) 30 minutes, room temperature
STEP(1.2) 1 hour, room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 15 hours, 70 deg C, 60 atm


RX(201) OF 350 - 3 STEPS



1.1. BuLi, THF
1.2. THF
2. Tri-*o*-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
3. Pd(OAc)₂,
1,10-Phenanthroline,
CO, DME

Print selected from 10562215.trn

RX(201) OF 350 - 3 STEPS




94%

NOTE: 2) stereoselective

CON: STEP(1.1) 30 minutes, room temperature
STEP(1.2) 1 hour, room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 14 hours, 70 deg C, 15 psi


RX(202) OF 350 - 3 STEPS



1.1. BuLi, THF
1.2. THF
2. Tri-*o*-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
3. Pd(OAc)₂, PPh₃, CO,
MeCN

Print selected from 10562215.trn


RX(202) OF 350 - 3 STEPS



NOTE: 2) stereoselective

CON: STEP(1.1) 30 minutes, room temperature
STEP(1.2) 1 hour, room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 15 hours, 70 deg C, 60 atm


RX(206) OF 350 - 4 STEPS



1. MeOH, KOH
- 2.1. Ph₃PMe.Br, BuLi,
THF
- 2.2. THF
3. Tri-*o*-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
4. Pd(OAc)₂,
1,10-Phenanthroline,
CO, DMF

Print selected from 10562215.trn


RX(206) OF 350 - 4 STEPS



NOTE: 3) stereoselective

CON: STEP(1) 2.5 hours, reflux
STEP(2.1) 30 minutes, room temperature
STEP(2.2) 1 hour, room temperature
STEP(3) 4 hours, 100 deg C
STEP(4) 14 hours, 70 deg C, 15 psi



RX(207) OF 350 - 4 STEPS



1. MeOH, KOH
- 2.1. Ph₃PMe.Br, BuLi,
THF
- 2.2. THF
3. Tri-*o*-tolylphosphine,
Pd(OAc)₂, Et₃N,
DMF
4. Pd(OAc)₂, PPh₃, CO,
MeCN

Print selected from 10562215.trn

RX(207) OF 350 - 4 STEPS





NOTE: 3) stereoselective

CON: STEP(1) 2.5 hours, reflux
STEP(2.1) 30 minutes, room temperature
STEP(2.2) 1 hour, room temperature
STEP(3) 4 hours, 100 deg C
STEP(4) 15 hours, 70 deg C, 60 atm

Print selected from 10562215.trn



RX(214) OF 350 - 5 STEPS



NOTE: 4) stereoselective


CON: STEP(1.1) 0 deg C, 2 hours, room temperature
STEP(2) 4 hours, room temperature
STEP(3) 4 hours, room temperature
STEP(4) 4 hours, 100 deg C
STEP(5) 14 hours, 70 deg C, 15 psi

RX(215) OF 350 - 5 STEPS



Print selected from 10562215.trn

RX (215) OF 350 - 5 STEPS



NOTE: 4) stereoselective

CON: STEP(1.1) 0 deg C; 2 hours, room temperature



STEP(2) 4 hours, room temperature

STEP(3) 4 hours, room temperature

STEP(4) 4 hours, 100 deg C

STEP(5) 15 hours, 70 deg C, 60 atm

RX (220) OF 350 - 5 STEPS



94%

NOTE: 3) key intermediate, 4) stereoselective

CON: STEP(1.1) 0 deg C; 2 hours, room temperature

STEP(2.1) -20 deg C; <-5 deg C; 30 minutes, -20 deg C

STEP(2.2) 3 hours, room temperature

STEP(3.1) room temperature; 30 minutes, room temperature



STEP(4.1) 30 minutes, room temperature

STEP(4.2) 30 minutes, 60 deg C

STEP(5) 14 hours, 70 deg C, 15 psi


Print selected from 10562215.trn

RX (221) OF 350 - 5 STEPS



95%

RX (221) OF 350 - 5 STEPS



NOTE: 3) key intermediate, 4) stereoselective

CON: STEP(1.1) 0 deg C; 2 hours, room temperature

STEP(2.1) -20 deg C; <-5 deg C; 30 minutes, -20 deg C

STEP(2.2) 3 hours, room temperature

STEP(3.1) room temperature; 30 minutes, room temperature


STEP(4.1) 30 minutes, room temperature

STEP(4.2) 30 minutes, 60 deg C

STEP(5) 15 hours, 70 deg C, 60 atm

Print selected from 10562215.trn


RX (226) OF 350 - 5 STEPS



NOTE: 4) stereoselective


CON: STEP(1.1) 20 minutes, 0 deg C; 30 minutes, room temperature
 STEP(1.2) 30 minutes, room temperature
 STEP(2) 4 hours, room temperature
 STEP(3) 4 hours, room temperature
 STEP(4) 4 hours, 100 deg C
 STEP(5) 14 hours, 70 deg C, 15 psi

RX (227) OF 350 - 5 STEPS



Print selected from 10562215.trn


RX (227) OF 350 - 5 STEPS



NOTE: 4) stereoselective

CON: STEP(1.1) 20 minutes, 0 deg C; 30 minutes, room temperature
 STEP(1.2) 30 minutes, room temperature
 STEP(2) 4 hours, room temperature
 STEP(3) 4 hours, room temperature
 STEP(4) 4 hours, 100 deg C
 STEP(5) 15 hours, 70 deg C, 60 atm

RX (232) OF 350 - 5 STEPS




NOTE: 3) key intermediate, 4) stereoselective


CON: STEP(1.1) 20 minutes, 0 deg C; 30 minutes, room temperature
 STEP(1.2) 30 minutes, room temperature
 STEP(2.1) -20 deg C; <-5 deg C; 30 minutes, -20 deg C
 STEP(2.2) 3 hours, room temperature
 STEP(3.1) room temperature; 30 minutes, room temperature
 STEP(4.1) 30 minutes, room temperature
 STEP(4.2) 30 minutes, 60 deg C
 STEP(5) 14 hours, 70 deg C, 15 psi

Print selected from 10562215.trn

RX(233) OF 350 - 5 STEPS



RX(233) OF 350 - 5 STEPS




NOTE: 3) key intermediate, 4) stereoselective

CON: STEP(1.1) 20 minutes, 0 deg C; 30 minutes, room temperature
 STEP(1.2) 30 minutes, room temperature
 STEP(2.1) -20 deg C; <-5 deg C; 30 minutes, -20 deg C
 STEP(2.2) 3 hours, room temperature
 STEP(3.1) room temperature; 30 minutes, room temperature
 STEP(4.1) 30 minutes, room temperature
 STEP(4.2) 30 minutes, 60 deg C
 STEP(5) 15 hours, 70 deg C, 60 atm

Print selected from 10562215.trn

RX(234) OF 350 - 5 STEPS




HCl
100%

NOTE: 3) stereoselective

CON: STEP(1) 4 hours, room temperature
 STEP(2) 4 hours, room temperature
 STEP(3) 4 hours, 100 deg C
 STEP(4) 14 hours, 70 deg C, 15 psi


RX(235) OF 350 - 5 STEPS



1. Pd, H2, AcOEt
 2. C:98327-87-8,
 R:338746-12-6, KBr,
 Water, AcOH
 3. Tri-o-tolylphosphine,
 Pd(OAc)2, Et3N,
 DMF
 4. Pd(OAc)2, PPh3, CO,
 MeCN
 5. HCl, Water, DMF

Print selected from 10562215.trn

RX(235) OF 350 - 5 STEPS





HCl
100%

NOTE: 3) stereoselective

CON: STEP(1) 4 hours, room temperature
STEP(2) 4 hours, room temperature
STEP(3) 4 hours, 100 deg C
STEP(4) 15 hours, 70 deg C, 60 atm

RX(237) OF 350 - 6 STEPS





HCl
100%

NOTE: 4) stereoselective

CON: STEP(1.1) 0 deg C; 2 hours, room temperature
STEP(2) 4 hours, room temperature
STEP(3) 4 hours, room temperature
STEP(4) 4 hours, 100 deg C
STEP(5) 14 hours, 70 deg C, 15 psi

Print selected from 10562215.trn

RX(238) OF 350 - 6 STEPS




HCl
100%

NOTE: 4) stereoselective

CON: STEP(1.1) 0 deg C; 2 hours, room temperature
STEP(2) 4 hours, room temperature
STEP(3) 4 hours, room temperature
STEP(4) 4 hours, 100 deg C
STEP(5) 15 hours, 70 deg C, 60 atm

Print selected from 10562215.trn

RX(240) OF 350 - 6 STEPS




HCl
100%

NOTE: 4) stereoselective

CON: STEP(1.1) 20 minutes, 0 deg C; 30 minutes, room temperature
STEP(1.2) 30 minutes, room temperature
STEP(2) 4 hours, room temperature
STEP(3) 4 hours, room temperature
STEP(4) 4 hours, 100 deg C
STEP(5) 14 hours, 70 deg C, 15 psi

RX(241) OF 350 - 6 STEPS




HCl
100%

Print selected from 10562215.trn

NOTE: 4) stereoselective


CON: STEP(1.1) 20 minutes, 0 deg C; 30 minutes, room temperature
STEP(1.2) 30 minutes, room temperature
STEP(2) 4 hours, room temperature
STEP(3) 4 hours, room temperature
STEP(4) 4 hours, 100 deg C
STEP(5) 15 hours, 70 deg C, 60 atm

RX(244) OF 350 - 5 STEPS



converging
Ph3CMe.Br

RX(244) OF 350 - 5 STEPS




HCl
100%

NOTE: stereoselective


CON: STEP(1.1) 4 hours, room temperature
STEP(1.2) room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 14 hours, 70 deg C, 15 psi
STEP(5.1) 30 minutes, room temperature
STEP(5.2) 1 hour, room temperature

Print selected from 10562215.trn

RX(245) OF 350 - 5 STEPS



RX(245) OF 350 - 5 STEPS




HCl
100%

NOTE: stereoselective


CON: STEP(1.1) 4 hours, room temperature
STEP(1.2) room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 15 hours, 70 deg C, 60 atm
STEP(5.1) 30 minutes, room temperature
STEP(5.2) 1 hour, room temperature

Print selected from 10562215.trn

RX(247) OF 350 - 4 STEPS



RX(247) OF 350 - 4 STEPS




94%

NOTE: stereoselective



CON: STEP(1.1) 4 hours, room temperature
STEP(1.2) room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 14 hours, 70 deg C, 15 psi
STEP(4.1) 30 minutes, room temperature
STEP(4.2) 1 hour, room temperature

RX(248) OF 350 - 4 STEPS



Print selected from 10562215.trn


RX(248) OF 350 - 4 STEPS



NOTE: stereoselective


CON: STEP(1.1) 4 hours, room temperature
STEP(1.2) room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 15 hours, 70 deg C, 60 atm
STEP(4.1) 30 minutes, room temperature
STEP(4.2) 1 hour, room temperature

RX(250) OF 350 - 5 STEPS



Print selected from 10562215.trn


RX(250) OF 350 - 5 STEPS



NOTE: stereoselective



CON: STEP(1.1) 4 hours, room temperature
STEP(1.2) room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 14 hours, 70 deg C, 15 psi
STEP(4) 2.5 hours, reflux
STEP(5.1) 30 minutes, room temperature
STEP(5.2) 1 hour, room temperature

RX(251) OF 350 - 5 STEPS



Print selected from 10562215.trn



RX(251) OF 350 - 5 STEPS



NOTE: stereoselective


CON: STEP(1.1) 4 hours, room temperature
 STEP(1.2) room temperature
 STEP(2) 4 hours, 100 deg C
 STEP(3) 15 hours, 70 deg C, 60 atm
 STEP(4) 2.5 hours, reflux
 STEP(5.1) 30 minutes, room temperature
 STEP(5.2) 1 hour, room temperature

RX(253) OF 350 - 5 STEPS



Print selected from 10562215.trn

RX(253) OF 350 - 5 STEPS





HCl
100%

NOTE: stereoselective


CON: STEP(1) 4 hours, room temperature
 STEP(2) 4 hours, 100 deg C
 STEP(3) 14 hours, 70 deg C, 15 psi
 STEP(5.1) 30 minutes, room temperature
 STEP(5.2) 1 hour, room temperature

RX(254) OF 350 - 5 STEPS



Print selected from 10562215.trn



RX(254) CF 350 - 5 STEPS




941
100%

NOTE: stereoselective
CON: STEP(1) 4 hours, room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 15 hours, 70 deg C, 60 atm
STEP(5.1) 30 minutes, room temperature
STEP(5.2) 1 hour, room temperature

RX(256) CF 350 - 4 STEPS



converging,
Ph3PMe.Br





944

<-----User Break----->


NOTE: stereoselective
CON: STEP(1) 4 hours, room temperature
STEP(2) 4 hours, 100 deg C
STEP(3) 14 hours, 70 deg C, 15 psi
STEP(4.1) 30 minutes, room temperature
STEP(4.2) 1 hour, room temperature

Print selected from 10562215.trn

RX(257) CF 350 - 4 STEPS



converging
Ph3PMe.Br
Ph3PMe.Br





954

=> d chib abs ford 1-
YOU HAVE REQUESTED DATA FROM 2 ANSWERS - CONTINUE? Y/N:y

LB ANSWER 1 OF 2 CASREACT COPYRIGHT 2009 ACS on STN
144:292572 Diaminocoumarins as fluorogenic substrates for monoamine oxidases,
and their preparation, pharmaceutical compositions, photophysical
properties, and a method for detecting active monoamine oxidases and their
inhibitors for treatment of nervous system disorders. Chen, Gong; Yee,
Dominic J.; Gubernator, Nikolai; Sames, Dalibor (The Trustees of Columbia
University in the City of New York, USA). PCT Int. Appl. WO 2006026368
A2 20060909, 278 pp. DESIGNATED STATES: K: AE, AG, AL, AM, AT, AU, AZ,
BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, DE, DK, DM, DO,
EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HD, ID, IL, IN, IS, JP, KE,
KG, KM, KN, KR, KZ, LC, LE, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, ME, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,
SK, SL, SM, ST, TJ, TK, TN, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA,
ZM, ZW; RW: AT, BE, BF, BG, CH, CG, CI, CM, CY, CZ, DE, ES, FI, FR,
GA, GB, GR, IE, IS, IT, LU, MC, ML, MR, NE, NG, PT, SE, SN, TD, TG, TR.
(English). CODEN: FIMXOZ. APPLICATION: WO 2005-0830276 20050825.
PRIORITY: US 2004-64558P 20040825.

GI

Print selected from 10562215.tn




AB The invention relates to compds. of formula I, which are useful for detecting the activity of monoamine oxidases (MAO), compds. useful for competitively inhibiting monoamine oxidases, for determining inhibitors of monoamine oxidases and compds. useful for treating monoamine oxidase-related nervous system pathologies, as well as pharmaceutical compds. and methods of manufacture thereof. Compds. of formula I wherein R1 is H, alkyl, alkenyl, alkynyl, (un)substituted (hetero)aryl, cycloalkyl, NH2 and derivs., alkyl-CO2H, alkyl-OH, alkyl-NH2, halo, CX3, or indole; R2-R6 are independently H, OH, alkyl, alkenyl, alkynyl, (un)substituted (hetero)aryl, cycloalkyl, NH2 and derivs., alkyl-CO2H, alkyl-OH, alkyl-NH2, O-alkyl, O-alkenyl, O-alkynyl, O-aryl, O-cycloalkyl, CX3, halo, or indole; R1R2 and R1R6 may independently form an unsubstituted pyrrole; R1R2R6 may form an octahydroquinazolinone; R2R3 may form a pyrrole; X is halo; or pharmaceutically acceptable salts, or stereoisomers thereof are claimed in this invention. The process for preparing these fluorogenic substrates, and a method for identifying a test compound as a substrate of MAO are also claimed. Example compound II was prepared by nitration of 7-methylcoumarin, and the resulting 7-methyl-6-nitrocoumarin underwent condensation with DMF di-Me acetal to give the corresponding coumarin-enamine, which was hydrolyzed to give 7-(2-oxoethyl)-6-nitrocoumarin, which underwent reductive amination with dibenzosuberylaminoethylamine; the resulting 7-(dibenzosuberylaminoethyl)-6-nitrocoumarin was reduced to the 6-amino coumarin compound, which was hydrolyzed to give coumarin II. The invention compds. were evaluated for their photophys. properties and their enzymic activity toward MAO-A and MAO-B. Coumarin II showed λ_{max} 352 nm, ϵ 2000-100 M⁻¹cm⁻¹, ϕ 0.017 \pm 0.001, and λ_{em} was not detected. The invention compds. were evaluated for their ability to convert to the corresponding indole in the presence of MAO-A and MAO-B. Enzyme-catalyzed indole formation was realized after 24 h by reading fluorescence emission upon excitation at the absorbance maxima of the resp. indoles. Only diamino-coumarin II showed a significant conversion to its indole III in the presence of MAO-A and MAO-B. The enzyme kinetics for fluorogenic probe II were also determined. Compound II showed for MAO-A: K_m = 30.3 \pm 1.8 μ M, K_{cat} = 0.475 min⁻¹, K_{cat}/K_m = 0.015 min⁻¹ μ M⁻¹, V_{max} (mitochondria) = 0.0570 nmol min⁻¹ mg⁻¹; and for MAO-B: K_m = 51 \pm 140 μ M, K_{cat} = 20.62 min⁻¹, K_{cat}/K_m = 0.040 min⁻¹ μ M⁻¹, and V_{max} (mitochondria) = 1.2 nmol min⁻¹ mg⁻¹. These results indicate that coumarin probe II is a good substrate for monoamine oxidase and could be useful as a fluorescent reporter for monoamine

Print selected from 10562215.tn

oxidase.

RX(15) OF 512



CON: 2 hours, room temperature


18 ANSWER 2 OF 2 CASREACT COPYRIGHT 2009 ACS on STN
142:41180 Synthesis of 5-Substituted-1H-indol-2-yl-1H-guinin-2-ones: A Novel Class of KDR Kinase Inhibitors. Kuethe, Jeffrey T.; Wong, Audrey; Qu, Chuanxing; Smitrovich, Jacqueline; Davies, Ian W.; Hughes, David L. (Department of Process Research, Merck & Co., Inc., Rahway, NJ, 07065, USA). Journal of Organic Chemistry, 70(7), 2555-2567 (English) 2005. CODEN: JOCEAH. ISSN: 0022-3263. Publisher: American Chemical Society.

GI



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB A number of approaches for the synthesis of the 1H-indol-2-yl-1H-guinin-2-one ring system found in the potent and selective KDR kinase inhibitor I are described. The preparation and reaction of trimethylsilylnitrobenzene II with 2-methoxy-3-quinolinecarboxaldehyde afforded alc. III, which was the key intermediate for the preparation of the target compounds. Conversion of alc. III to either nitroketone IV or nitrostyrene V set the stage for reductive cyclization. The quinolin-2-one functionality was unmasked in the last step to provide compound I in 56-60% overall yield from readily available starting materials.

RX(22) OF 350




Print selected from 10562215.trn



NOTE: Raney Nickel used
CON: 7.5 hours, 65 deg C, 40 psi

=>
Uploading C:\Program Files\Straxp\Queries\10597127-99.str




chain nodes :
7 8 9
ring nodes :
1 2 3 4 5 6 11 12 13 14 15 16 17 18 19
ring/chain nodes :
10
chain bonds :
5-8 6-7 8-9 9-10
ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6 11-12 11-16 12-13 13-14 14-15 15-16 15-17 16-19
17-18 18-19
exact/norm bonds :
9-10 15-17 16-19 17-18 18-19
exact bonds :
5-8 6-7 8-9
normalized bonds :
1-2 1-6 2-3 3-4 4-5 5-6 11-12 11-16 12-13 13-14 14-15 15-16

Print selected from 10562215.trn

Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS 9:CLASS 10:CLASS
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:Atom
fragments assigned product role:
containing 11
fragments assigned reactant/reagent role:
containing 1
node mappings:
6:16 6:16

L9 STRUCTURE UPLOADED

=> d 19
L9 HAS NO ANSWERS
L9 STR



Structure attributes must be viewed using STN Express query preparation.

=> s 19
SAMPLE SEARCH INITIATED 11:21:40 FILE 'CASREACT'
SCREENING COMPLETE - 268 REACTIONS TO VERIFY FROM 26 DOCUMENTS
100.0% DONE 268 VERIFIED 76 HIT RXNS 6 DOCS
SEARCH TIME: 00.00.01
FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED VERIFICATIONS: 4378 TO 6342
PROJECTED ANSWERS: 6 TO 266
L10 6 SEA SSS SM L9 (76 REACTIONS)
=> s 19 full
FULL SEARCH INITIATED 11:21:48 FILE 'CASREACT'
SCREENING COMPLETE - 6095 REACTIONS TO VERIFY FROM 502 DOCUMENTS
100.0% DONE 6095 VERIFIED 1697 HIT RXNS (120 INCOMP) 151 DOCS
SEARCH TIME: 00.00.02
L11 151 SEA SSS FUL L9 (1697 REACTIONS)
=> d his

Print selected from 10562215.trn

(FILE 'HOME' ENTERED AT 11:15:49 ON 16 MAR 2009)
FILE 'CASREACT' ENTERED AT 11:16:04 ON 16 MAR 2009
11 STRUCTURE UPLOADED
12 3 S L1
13 114 S L1 FULL
14 0 S L3 AND CARBON MONOXIDE
15 5849 S CARBON MONOXIDE

FILE 'REGISTRY' ENTERED AT 11:17:25 ON 16 MAR 2009
16 1 S CARBON MONOXIDE/CN

FILE 'CASREACT' ENTERED AT 11:17:52 ON 16 MAR 2009
17 10878 S 630-08-0
18 2 S L7 AND L3
19 STRUCTURE UPLOADED
20 6 S L9
21 151 S L9 FULL

=> s l11 and l7
22 3 L11 AND L7

=> s l12 not l8
23 1 L12 NOT L8

=> d cblb abs ford

24 ANSWER 1 OF 1 CASREACT COPYRIGHT 2009 ACS on STM
149:79436 Palladium-catalyzed synthesis of 3-indolecarboxylic acid
derivatives. Soderberg, Bjorn C. G.; Banini, Serge R.; Turrer, Michael
R.; Minter, Aaron B.; Arrington, Amanda K. (C. Eugene Bennett Department
of Chemistry, West Virginia University, Morgantown, WV, 26506-6045, USA).
Synthesis (6), 903-912 (English) 2008. CODEN: SYNTEP. ISSN: 0039-7881.
Publisher: Georg Thieme Verlag.


GI



AB Indoles having an ester functionality in the 3-position, e.g., I, were prepared from 2-(2-nitrophenyl)propenoic acid derivs. via a palladium-catalyzed reductive N-heteroannulation using carbon monoxide as the ultimate reducing agent. The starting materials were prepared either by a Stille coupling of 2-halo-1-nitrobenzenes with Et 2-(tributylstannyl)-2-propenoate or by vicarious nucleophilic substitution of nitrobenzenes followed by a Knoevenagel-type condensation with an aldehyde. Synthesis of an example of a 3-nitrile- and a 3-sulfone-substituted indole is also described using the same type of methodologies.

Print selected from 10562215.trn

RX(2) OF 53



CON: STAGE(1) room temperature; room temperature -> 110 deg C, 6 atm;
72 hours, 110 deg C, 6 atm

=> d his

(FILE 'HOME' ENTERED AT 11:15:49 ON 16 MAR 2009)
FILE 'CASREACT' ENTERED AT 11:16:04 ON 16 MAR 2009
25 STRUCTURE UPLOADED
26 3 S L1
27 114 S L1 FULL
28 0 S L3 AND CARBON MONOXIDE
29 5849 S CARBON MONOXIDE

FILE 'REGISTRY' ENTERED AT 11:17:25 ON 16 MAR 2009
30 1 S CARBON MONOXIDE/CN

FILE 'CASREACT' ENTERED AT 11:17:52 ON 16 MAR 2009
31 10878 S 630-08-0
32 2 S L7 AND L3
33 STRUCTURE UPLOADED
34 6 S L9
35 151 S L9 FULL
36 3 S L11 AND L7
37 1 S L12 NOT L8

=> s help
Enter the terms you wish to use in your search profile.
ENTER LOGIC EXPRESSION, QUERY NAME, OR (END):end
SEARCH ENDED BY USER

=> help sfields

The searchable fields in the CASREACT File are listed below. If you do not specify a field, your term will be searched in the Basic Index, which contains CAS Registry Numbers (R) for reactants, products, reagents, catalysts, and solvents, and single words from the reaction notes, title, supplementary terms, index terms, and abstracts. Simultaneous left and right truncation (SLART), e.g., S 770XINZ, may be used for search terms in the /BI and /NTE search fields in CASREACT.

Reaction Fields

Definition

Search Code

Print selected from 10562215.trn

-----	-----
Basic Index	/BI or none
Catalyst	/CAT
Functional Group Formed	/FG.FORM
Functional Group in Product	/FG.PRO
Functional Group in Reactant	/FG.RCT
Functional Group in Reactant, Reagent, or Product	/FG
Functional Group in Reagent	/FG.RGT
Functional Group Reacting	/FG.RXN
Functional Group Yield	/FG.YD
Functional Group Yield Data	/FG.YDT
NonProduct	/NPRO
Nonreacting Functional Group	/FS.NON
Number of Steps	/NS
Product	/PRO
Reactant	/RCT
Reactant or Reagent	/RRT
Reaction Notes	/RTE
Reagent	/RGT
Solvent	/SOL
Yield	/YD
Yield Data	/YDT

General Document Search Fields

Definition	Search Code
-----	-----
Basic Index	/BI
Abstract	/AB
Accession Number	/AN
Author or Inventor	/AU
CA Section Cross-reference	/SX
Classification Code (CA Section)	/CC
Controlled Term	/CT
Controlled Word	/CW
Corporate Source	/CS
Country of Author	/CTA
Crossover Key	/CK
Document Type	/DT
Entry Date	/ED
Field Availability	/FA
File Segment	/FS
Index Term	/IT
International Standard (Document) Number	/ISN
Issue Number of Publication	/IS
Journal Title	/JT
Language	/LA
Other Source	/OS
Publication Date	/PD
Publication Year	/PY
Publisher	/PB
Publisher Item Identifier	/PUI
Source	/SO
Supplementary Term	/ST
Title	/TI
Uniform Resource Locator	/URL

Print selected from 10562215.trn

Update Date	/UP
Volume and Issue of CA	/VI
Volume Number of Publication	/VL
Patent Search Fields	
Definition	Search Code
-----	-----
Designated States	/DS
International Patent Classification (includes Main and Secondary IPC)	/IC
International Patent Classification, Additional or Supplementary	/ICA
International Patent Classification, Index or Complementary	/ICI
International Patent Classification, Main	/ICM
International Patent Classification, Main Group, Range Searchable	/MGR
International Patent Classification, Secondary	/ICS
International Patent Classification, Subgroup, Range Searchable	/SGR
Inventor Name	/IN
National Patent Classification	/NCL
National Patent Classification, Range Searchable	/NCLR
Patent Application Country	/AC
Patent Application Date	/AD
Patent Application Number	/AP
Patent Application Year	/AY
Patent Assignee	/PA
Patent Country	/PC
Patent Kind Code	/PK
Patent Number	/PN
Priority Application Country	/PRC
Priority Application Date	/PRD
Priority Application Number	/PRN
Priority Application Year	/PRY

You may also use the following super search fields to execute a search in one or more patent fields:

Search Field Name	Super Search Code	Fields Searched
International Patent Classifications	/IPC	/IC, /ICA, /ICI
Patent Application and Priority Number	/APPS	/AP, /PRN
Patent Countries	/PCS	/PC, /DS
Patent Numbers	/PATIS	/PN

All fields are text fields except Entry Date (/ED), Functional Group Yield (/FG.YD), International Patent Classification, Main Group, Range Searchable (/MGR), International Patent Classification, Subgroup, Range Searchable (/SGR), Issue Number (/IS), National Patent Classification, Range Searchable (/NCLR), Number of Steps (/NS), Patent Application Date (/AD), Patent Application Year (/AY), Priority Application Date (/PRD), Priority Application Year (/PRY),

Print selected from 10562215.trn

Publication Date (/PD), Publication Year (/PY), Update Date (/UP), Volume (/VL) and Yield (/YD), which are numeric and may be searched with numeric operators or ranges, e.g., ED>19971230 or 1/NS.

Search and display fields generally have the same field codes. To see a list of display fields, enter HELP DFIELDS at an arrow prompt (=>).

=> d his
'HIA' IS NOT A VALID FORMAT FOR FILE 'CASREACT'

The following are valid formats:

ABS ----- GI and AB
ALL ----- BIB, AB, IND, RE, Single-step Reactions
APPS ----- AT, FRAI
BIB ----- AV, plus Bibliographic Data
CAN ----- List of CA abstract numbers without answer numbers
CBIB ----- AV, plus Compressed Bibliographic Data
DALL ----- ALL, delimited (end of each field identified)
IABS ----- ABS, indented with text labels
IALL ----- ALL, indented with text labels
IBIB ----- BIB, indented with text labels
IND ----- Indexing data
IPC ----- International Patent Classifications
ISTD ----- STD, indented with text labels
OIBIB ----- AV, plus Bibliographic Data (original)
OBIB ----- OBIB, indented with text labels

SRIB ----- BIB, no citations
SIBIB ----- IBIB, no citations

MAX ----- Same as ALL
PATS ----- PI, SO
SCAN ----- TI and FCRD (random display, no answer number. SCAN must be entered on the same line as DISPLAY, e.g., D SCAN.)
SSRX ----- Single-Step Reactions (Map, Diagram, and Summary for all single-step reactions)
STD ----- BIB, IPC, and NCL

CRD ----- Compact Display of All Hit Reactions
CRDREF ----- Compact Reaction Display and SO, PY for Reference
FHIT ----- Reaction Map, Diagram, and Summary for first hit reaction
FHITCBIB ----- FHIT, AV plus CBIB
FCRD ----- First hit in Compact Reaction Display (CRD) format
FCRDREF ----- First hit in Compact Reaction Display (CRD) format with CA reference information (SO, PY). (Default)
FPAH ----- PATH, plus Reaction Summary for the "long path"
FSPATH ----- SPATH, plus Reaction Summary for the "short path"
HIT ----- Reaction Map, Reaction Diagram, and Reaction Summary for all hit reactions and fields containing hit terms
OCC ----- All hit fields and the number of occurrences of the hit terms in each field. Includes total number of HIT, PATH, SPATH reactions. Labels reactions that have incomplete verifications.

Print selected from 10562215.trn

PATH ----- Reaction Map and Reaction Diagram for the "long path". Displays all hit reactions, except those whose steps are totally included within another hit reaction which is displayed
RX ----- Hit Reactions (Map, Diagram, Summary for all hit reactions)
RXG ----- Hit Reaction Graphics (Map and Diagram for all hit reactions)
RXL ----- Hit Reaction Long (Map, Diagram, Summary for all hit reactions)
RXS ----- Hit Reaction Summaries (Map and Summary for all hit reactions)
SPATH ----- Reaction Map and Reaction Diagram for the "short path". Displays all single step reactions which contain a hit substance. Also displays those multistep reactions that have a hit substance in both the first and last steps of the reaction, except for those hit reactions whose steps are totally included within another hit reaction which is displayed

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of combinations include: D TI; D BIB RXG; D TI, AV, FCRD. The information is displayed in the same order as the specification. All of the formats, except CRD, CRDREF, FHIT, PATH, FPAH, SPATH, FSPATH, FCRD, FCRDREF, HIT, RX, RXG, RXS, SCAN, and OCC, may be used with the DISPLAY command to display the record for a specified Accession Number.

ENTER DISPLAY FORMAT (FCRDREF):d his
'D' IS NOT A VALID FORMAT FOR FILE 'CASREACT'

The following are valid formats:

ABS ----- GI and AB
ALL ----- BIB, AB, IND, RE, Single-step Reactions
APPS ----- AT, FRAI
BIB ----- AV, plus Bibliographic Data
CAN ----- List of CA abstract numbers without answer numbers
CBIB ----- AV, plus Compressed Bibliographic Data
DALL ----- ALL, delimited (end of each field identified)
IABS ----- ABS, indented with text labels
IALL ----- ALL, indented with text labels
IBIB ----- BIB, indented with text labels
IND ----- Indexing data
IPC ----- International Patent Classifications
ISTD ----- STD, indented with text labels
OIBIB ----- AV, plus Bibliographic Data (original)
OBIB ----- OBIB, indented with text labels

SRIB ----- BIB, no citations
SIBIB ----- IBIB, no citations

MAX ----- Same as ALL
PATS ----- PI, SO
SCAN ----- TI and FCRD (random display, no answer number. SCAN must be entered on the same line as DISPLAY, e.g., D SCAN.)
SSRX ----- Single-Step Reactions (Map, Diagram, and Summary for all single-step reactions)
STD ----- BIB, IPC, and NCL

CRD ----- Compact Display of All Hit Reactions

Print selected from 10562215.trn

CRDREF ----- Compact Reaction Display and SO, PY for Reference
 FHIT ----- Reaction Map, Diagram, and Summary for first hit reaction
 FHITCBIB ---- FHIT, AM plus CBIB
 FCRO ----- First hit in Compact Reaction Display (CRO) format
 FCROREF ---- First hit in Compact Reaction Display (CRO) format with CA reference information (SO, PY). (Default)
 FPATH ----- PATH, plus Reaction Summary for the "long path"
 FSPATH ----- SPATH, plus Reaction Summary for the "short path"
 HIT ----- Reaction Map, Reaction Diagram, and Reaction Summary for all hit reactions and fields containing hit terms
 OCC ----- All hit fields and the number of occurrences of the hit terms in each field. Includes total number of HIT, PATH, SPATH reactions. Labels reactions that have incomplete verifications.
 PATH ----- Reaction Map and Reaction Diagram for the "long path". Displays all hit reactions, except those whose steps are totally included within another hit reaction which is displayed
 RX ----- Hit Reactions (Map, Diagram, Summary for all hit reactions)
 RXG ----- Hit Reaction Graphics (Map and Diagram for all hit reactions)
 RXL ----- Hit Reaction Long (Map, Diagram, Summary for all hit reactions)
 RXS ----- Hit Reaction Summaries (Map and Summary for all hit reactions)
 SPATH ----- Reaction Map and Reaction Diagram for the "short path". Displays all single step reactions which contain a hit substance. Also displays those multistep reactions that have a hit substance in both the first and last steps of the reaction, except for those hit reactions whose steps are totally included within another hit reaction which is displayed

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDOS at an arrow prompt (=>). Examples of combinations include: D TI; D BIR RX; D TI, AU, FCRO. The information is displayed in the same order as the specification. All of the formats, except CRD, CRDREF, FHIT, PATH, FPATH, SPATH, FSPATH, FCRO, FCROREF, HIT, RX, RXG, RXS, SCAN, and OCC, may be used with the DISPLAY command to display the record for a specified Accession Number.

ENTER DISPLAY FORMAT (FCROREF):end

=> d his

(FILE 'HWE' ENTERED AT 11:15:49 ON 16 MAR 2009)
 FILE 'CASREACT' ENTERED AT 11:16:04 ON 16 MAR 2009
 11 STRUCTURE UPLOADED
 12 3 S L1
 13 114 S L1 FULL
 14 4 S L3 AND CARBON MONOXIDE
 15 5849 S CARBON MONOXIDE
 FILE 'REGISTRY' ENTERED AT 11:17:25 ON 16 MAR 2009
 16 1 S CARBON MONOXIDE/CN
 FILE 'CASREACT' ENTERED AT 11:17:52 ON 16 MAR 2009
 17 10878 S 430-08-0

Print selected from 10562215.trn


L8 2 S L7 AND L3
 L9 STRUCTURE UPLOADED
 L10 6 S L9
 L11 151 S L9 FULL
 L12 3 S L11 AND L7
 L13 1 S L12 NOT L8
 => s l11 and iron
 15437 IRON
 12 IRONS
 15437 IRON
 (IRON OR IRONS)
 L14 3 L11 AND IRON
 => s l11 and carbonyl
 35779 CARBONYL
 1939 CARBONYLS
 36380 CARBONYL
 (CARBONYL OR CARBONYLS)
 L15 3 L11 AND CARBONYL
 => s l11 and complex
 100247 COMPLEX
 80122 COMPLEXES
 117248 COMPLEX
 (COMPLEX OR COMPLEXES)
 L16 2 L11 AND COMPLEX
 => s l13-116
 L17 9 (L13 OR L14 OR L15 OR L16)
 => s l14-116
 L18 8 (L14 OR L15 OR L16)
 => s l18 not l12
 L19 8 L18 NOT L12
 => s l18 not l8
 L20 8 L18 NOT L8
 => d cbib abs ford

L20 ANSWER 1 OF 8 CASREACT COPYRIGHT 2009 ACS on STM
 148:449603 Atom-efficient synthesis of 2,6-diazacyclophane compounds through alcoholysis/reduction of 3-nitroaryl-methylene-2,3-piperazinediones.
 Gonzalez, Juan Francisco; de la Cuesta, Elena; Avendano, Carmen
 (Departamento de Química Orgánica y Farmacéutica, Facultad de Farmacia, Universidad Complutense, Madrid, 28040, Spain). Tetrahedron, 64(12), 2762-2771 (English) 2008. CODEN: TETRAH. ISSN: 0040-4020. Publisher: Elsevier Ltd..
 AB A one-pot ring-opening/alcoholysis/hydrolysis process with 3-([nitroaryl]methylene)-2,3-piperazinediones yielded N-[3-([nitroaryl]pyrrolyl) amino esters, which gave the corresponding amines by reduction of the nitro group. In the case of 2-nitroaryl compds., an intramol. reductive amination afforded N-(indole-2-carbonyl) amino esters, while the intermol. reductive amination of 3- and 4-nitroaryl derivs. allowed the synthesis of 2,6-diazacyclophanes. The amine compds. may be coupled with amino acids to yield peptide-like

Print selected from 10562215.trn

derivs.

RX(22) OF 77



CON: 5 hours, room temperature

=> d cbb abs ford 1-

YOU HAVE REQUESTED DATA FROM 8 ANSWERS - CONTINUE? Y/N:y

L20 ANSWER 1 OF 8 CASREACT COPYRIGHT 2009 ACS on STM


148:449603 Atom-efficient synthesis of 2,6-diazacyclophane compounds through alcoholysis/reduction of 3-nitroarylmethylene-2,5-piperazinediones.

Gonzalez, Juan Francisco; de la Cuesta, Elena; Avendano, Carmen (Departamento de Química Orgánica y Farmacéutica, Facultad de Farmacia, Universidad Complutense, Madrid, 28040, Spain). *Tetrahedron*, 64(12), 2762-2771 (English) 2008. CODEN: TETRA3. ISSN: 0040-4020. Publisher: Elsevier Ltd..

AB A one-pot ring-opening/alcoholysis/hydrolysis process with 3-[(nitroaryl)methylene]-2,5-piperazinediones yielded N-[3-(nitroaryl)pyruvoyl] amino esters, which gave the corresponding amines by reduction of the nitro group. In the case of 2-nitroaryl compds., an intramol. reductive amination afforded N-(indole-2-carbonyl) amino esters, while the intermol. reductive amination of 3- and 4-nitroaryl derivs. allowed the synthesis of 2,6-diazacyclophanes. The amino compds. may be coupled with amino acids to yield peptide-like derivs.

Print selected from 10562215.trn

RX(22) OF 77



CON: 5 hours, room temperature


L20 ANSWER 2 OF 8 CASREACT COPYRIGHT 2009 ACS on STM

148:23738 Chemistry of Pyrrolo[1,2-a]indole- and Pyrido[1,2-a]indole-Based Quinone Methides. Mechanistic Explanations for Differences in

Cytostatic/Cytotoxic Properties. Khouri, Omar; Skibo, Edward B. (Department of Chemistry and Biochemistry, Arizona State University, Tempe, AZ, 85287-1604, USA). *Journal of Organic Chemistry*, 72(23), 8636-8647 (English) 2007. CODEN: JOCEAH. ISSN: 0022-3263. Publisher: American Chemical Society.

AB In the present study the authors investigate pyrrolo[1,2-a]indole- and pyrrolo[1,2-a]indole-based quinones capable of forming quinone methide and vinyl quinone species upon reduction and leaving group elimination. The goals were to determine the influence of the 6-membered pyrido and the 5-membered pyrrolo fused rings on quinone methide and vinyl quinone formation and fate as well as on cytostatic and cytotoxic activity. The authors used the technique of Spectral Global Fitting to study the fleeting quinone methide intermediate directly. Conclusions regarding quinone methide reactivity are that carbonyl O-protonation is required for nucleophile trapping and that the pK_a value of this protonated species is near neutrality. The abnormally high protonated carbonyl pK_a values are due to the formation of an aromatic carbocation species upon protonation. The fused pyrido ring promotes quinone methide and vinyl quinone formation but slows nucleophile trapping compared to the fused pyrrolo ring. These findings are explained by the presence of axial hydrogen atoms in the fused pyrido ring resulting in more steric congestion compared to the relatively flat fused pyrrolo ring. Consequently, pyrrolo[1,2-a]indole-based quinones exhibit more cytostatic activity than the pyrido[1,2-a]indole analogs due to their greater nucleophile trapping capability.

RX(11) OF 144



x K
(step 1)


CON: STAGE(1) room temperature; 4 hours, room temperature
STAGE(2) 3 hours, 40 - 45 deg C

L20 ANSWER 3 OF 8 CASREACT COPYRIGHT 2009 ACS on SIN

138:204905 2-Alkylindole-3-carboxylate esters by a tandem reduction-addition-elimination reaction. Bunce, Richard A.; Randall, Marty H.; Applegate, Kirby G. (Department of Chemistry, Oklahoma State University, Stillwater, OK, 74078-3071, USA). Organic Preparations and Procedures International, 34(5), 483-489 (English) 2002. CODEN: OPPIAR. ISSN: 0030-4948. Publisher: Organic Preparations and Procedures, Inc..

AB 2-Alkylindole-3-carboxylate esters were prepared by nucleophilic substitution of 1-fluoro-2-nitrobenzene with β -keto ester anions formed from $\text{R}^1\text{OCH}_2\text{CO}_2\text{Et}$ (R^1 = alkyl, 3-butenyl, $\text{CH}_2\text{CH}_2\text{Ph}$, cyclohexyl, Ph, 2-ClO $_2$ CH $_2$), to give (2)-(2-oxo-2-alkyl-1-oxoethyl)-3-(alkoxycarbonyl)indole, followed by tandem reduction-addition-elimination using iron powder in glacial acetic acid.

RX(11) OF 30




NOTE: Michael addn.

CON: STAGE(1) 115 deg C; 115 deg C -> room temperature

L20 ANSWER 4 OF 8 CASREACT COPYRIGHT 2009 ACS on SIN

133:266728 Preparation of N-hydroxypyrrole-2-carboxylic acids and pyrrole-2-carboxylic acids. Katayama, Seiji; Ae, Nobuyuki; Nagata, Tatsuo (Sumitomo Pharmaceuticals Co., Ltd., Japan). Jpn. Kokai Tokkyo Koho JP 2000273084 A 20001003, 13 pp. (Japanese). CODEN: JKKOAF. APPLICATION: JP 1999-77212 19990323.



61



AB N-hydroxypyrrole-2-carboxylic acids I [R^1 - R^3 = H, (un)substituted alkyl, (un)substituted aryl, (un)substituted heteroaryl; R^1 and R^2 may be bonded together to form a condensed aromatic ring; R^2 and R^3 may be bonded together to form a condensed cycloalkane or heterocycloalkane ring; R^4 = (un)substituted alkyl], useful as intermediates for drugs and agrochemicals, are prepared by treating nitro keto esters II (R^1 - R^4 = same as above) with reducing agents. Pyrrole-2-carboxylic acids III (R^1 - R^4 = same as above), also useful as intermediates for drugs and agrochemicals, are prepared by treating II with reducing agents and reacting the resulting I with other reducing agents. A dimethoxyethane solution of Me (2-iodo-4-chloro-6-nitrophenyl)pyruvate [preparation from 4-chloro-2-nitrotoluene (IV) with 2 steps] was gradually added to a H_2O /dimethoxyethane solution of SnCl_2 at room temperature and the reaction mixture was further stirred for 1 h. An aqueous TiCl_3 solution was gradually added to the above reaction mixture at 0° and the mixture was stirred for 1 h to give 51% (based on IV) Me 4-iodo-6-chloroindole-2-carboxylate. Preparation of 3-(S)-3-[2-(1-R-carboxylethoxy)-4-aminomethylphenyl]aminocarbonylmethyl-1,3,4,5-tetrahydrobenzo[d]indole-2-carboxylic acid hydrochloride monohydrate as a NMDA receptor antagonist.

Print selected from 10562215.trn

RX(1) OF 1




NOTE: room temp. for 1 h; ice-temp. for 1 h

L20 ANSWER 5 OF 8 CASREACT COPYRIGHT 2009 ACS on STN

104:109408 Synthesis of 4,7-indolequinones. The oxidative demethylation of 4,7-dimethoxyindoles with ceric ammonium nitrate. Kitahara, Yoshiyasu; Nakahara, Shinsuke; Murata, Ryuichi; Kubo, Akinori (Meiji Coll. Pharm., Tokyo, 154, Japan). Chemical & Pharmaceutical Bulletin, 33(5), 2122-8 (English) 1985. CODEN: CPBIAL. ISSN: 0009-2363.


GI



AB The indoles I (R, R3 = H, Me; R1 = Me, Et, Ph; R2 = H, OMe) were synthesized by reductive cyclization of the appropriate dinitrotyrosene or (nitrophenyl)acetaldehyde. The oxidative demethylations of I and phenol II with ceric ammonium nitrate yielded the corresponding 4,7-diones and p-quinone resp.

Print selected from 10562215.trn


RX(28) OF 151



L20 ANSWER 6 OF 8 CASREACT COPYRIGHT 2009 ACS on STN



89:4425 Indole derivatives. Batcho, Andrew David; Hengartner, Urs Oskar; Leingruber, Willy; Scott, John William; Valentine, Donald, Jr. (Hoffmann-La Roche, F., and Co. A.-G., Switz.). Ger. Offen. DE 2727671 19780105, 29 pp. (German). CODEN: GRXESX. APPLICATION: DE 1977-2727671 19770620.

GI



AB D-Tryptophans I (R and R1 = H, halo, OH, NH2, alkyl, alkoxy, aralkoxy, trihalomethyl; R2 = R3 = H) were prepared by asym. hydrogenating indoleacrylic acids II [R2 = alkyl, aryl; R3 = COR4 (R4 = H, alkyl, aryl, alkoxy, haloalkyl)] over a complex of a Rh compound and a chiral tertiary phosphine and hydrolyzing the resulting I [R2 = alkyl, aryl, ammonium, alkylammonium, R3 = COR4]. II were prepared by either condensing indolecarbaldehydes III with an alkyl acylammonium salt or by reducing the nitroacrylate IV (R5 = R2, R2 = same as in II) and acylating the resulting amine. Thus, 6-chloroindole was treated with EtOHC(CO2Me)CO2Me to give II.HCl (R = 6-Cl, R1 = R3 = H, R2 = Me), which was acetylated and saponified to give II (R = 6-Cl, R1 = R2 = H, R3 = Ac). The latter was hydrogenated over [Rh(1,5-cyclooctadiene)(Cl)]2 and chiro-trans-4,5-bis(di-n-tolylphosphino)methyl-2,2-dimethyl-1,3-dioxolane to give N-acetyl-6-chloro-D-tryptophan, which was hydrolyzed with 2N HBr to give 6-chloro-D-tryptophan.

RX (28) OF 69 - 2 STEPS



120 ANSWER 7 OF 8 CASREACT COPYRIGHT 2009 ACS on STN

50:27880 Ergot alkaloids. XL. A new synthesis of bufotenine and related hydroxytryptamines. Stoll, A.; Troxler, F.; Peyer, J.; Hofmann, A. (Sandoz, Basel, Switz.). *Helvetica Chimica Acta*, 38, 1452-72 (German) 1955. CODEN: HCACW. ISSN: 0018-019X.

AB cf. preceding abstract Nitrosation of *m*-MeC6H4OH and oxidation of the NO compound give 63% 2,5-(O2N)(HO)C6H3Me, m. 129-30°, which is converted into 87% 2,5-(O2N)(PhCH2O)C6H3Me (I). Treating I mole I with 2 mol (CO2Et)2 and 2 mol EtOK according to Burton and Stoves (C.A. 32, 550.1) at below 8° gives 87% 2-nitro-5-benzoyloxyphenylpyruvic acid, m. 112-13°, which (55 g.), reductively cyclized in 600 cc. H2O and 80 cc. 2N NaOH with 70 g. Na2S2O4 added in small portions until the color reaction (deep red) with NaOH is neg. and acidified with dilute HCl, gives 48.5% 5-benzoyloxyindole-2-carboxylic acid (II), m. 194-6°. Heating II in quinoline with Cu powder at 245-50° gives 80% 5-benzoyloxyindole (III), m. 103-5°, which, shaken in MeOH with Pd-asbestos (IV) and H, gives 5-hydroxyindole, long needles, m. 107-8°. Treating III in 1:1 EtOH-AcOH with Me2NH and CH2O according to Ek and Wiktor (C.A. 49, 12437i) gives 84% 5-benzoylgramine (V), m. 138°. Adding (20 min.) with stirring 420 cc. MeI to 30 g. V, heating the mixture 15 h. at 5°, keeping the methiodide with 60 g. MeCN in 1:1. H2O 2 h. at 80°, extracting the solution with CHCl3, evaporating the CHCl3, taking up the residue (29.6 g.) in 250 cc. Et2O, and diluting the concentrated Et2O solution with petr. ether give 65% 5-benzoyl-3-indoleacetoneitrile (VI), prisms, m. 75-8°. Refluxing 20 g. VI in 140 cc. EtOH and 100 cc. H2O 15 h. with 45 g. KOH, acidifying the mixture with 60 cc. AcOH, and diluting the filtered solution with 500 cc.

H2O



give 20.6 g. 5-benzoyl-3-indoleacetic acid, m. 145-7°, which is converted with CH2N2 into the Me ester and the latter heated with N2H4 1.5 h. at 135°, giving 95% 5-benzoyl-3-indoleacetylhydrazide (VII), leaflets, m. 153-4°. Adding dropwise 60 cc. N HCl to a mixture of 14.7 g. VII in 250 cc. dioxane and 50 cc. N NaNO2 solution, extracting the acetozide with Et2O, evaporating the Et2O, and treating the residual azide with 50 g. anhydrous Me2NH 3 h. at 5° give 60% 5-benzoyl-3-indoleacetodimethylamide (VIII), platelets, m. 138-40°. In a similar way the following admi. amides are prepared:

Me, short prisms, m. 141-2°; Et, prisms, m. 126-8°, d-l-Et, needles, m. 120-1°; H2NCH2CH2, plates, m. 137-9°; and piperidide, leaflets, m. 129-30°. Adding dropwise 1.26 g. LiAlH4 in 200 cc. Et2O in a N arm. to 3.65 g. VIII in 80 cc. THF, stirring the mixture 1 h. at 55°, and working it up in the usual way give 80% 5-benzoyloxy-*o*-N,N-dimethyltryptamine (bufotenine benzyl ether) (IX), pointed prisms, m. 87-9° [acid oxalate (X), fine leaflets, m. 177-8°]. Similar reduction of the corresponding amides gives the following *N*-substituted tryptamines: Me, plates, m. 84-6° [acid oxalate (XI), needles, m. 201-3°]; Et, crystals, m. 59-61° [acid oxalate, short needles, m. 187-9°] [the *o*-N,N-diethyl homolog does not crystallize (acid oxalate, prisms, m. 162°)]; H2NCH2CH2, does not crystallize (bis-acid oxalate, leaflets, m. 221-2°); *N*-(*β*-(5-benzoyl-3-indolyl)ethyl)piperidine, prisms, m. 136-8°. Shaking 3.45 g. IX in 75 cc. MeOH with 2 g. 5% IV and H 1.5 h. gives 78% bufotenine (XII), stout prisms, m. 138-40°. With FeCl3 in AcOH and concentrated H2SO4, XII gives a reddish color, turning to blue

after 1-2 s. The UV absorption curves of XII in EtOH, 0.1N HCl, and 0.1N NaOH, and the IR absorption curves of XII and of natural XII are given. Shaking 1.85 g. X in 200 cc. MeOH with IV in H gives 86% XII acid oxalate, needles, m. 89-90°. Treating 1.1 g. XII in 2 cc. MeOH with 2 cc. MeI 3 h. at 20° gives 1.7 g. XII methiodide, stout prisms, m. 214-15°. Dissolving 2.9 g. XII and 2.3 g. creatinine sulfate (XIII) in 14 cc. N H2SO4 and 40 cc. boiling H2O and diluting the solution with Me2CO give a 5:3 g. XII-XIII complex, fine needles, m. 147-9°. Debenzoylation of XI gives 5-hydroxy-*o*-N-methyltryptamine (*o*-N-methylserotonin), short pointed prisms and plates, m. 153-6°; *N*-Et homolog, short prisms, m. 239-40°; *N*,*N*-di-Et homolog, polyhedrons and prisms, m. 147-9° (oxalate, m. 230-2°); *N*-H2NCH2CH2 analog, bis-acid oxalate, leaflets, m. 208-9°; *N*-(*β*-(5-hydroxy-3-indolyl)ethyl)piperidine, stout prisms, m. 201-3° (oxalate, pointed prisms, m. 243-7°). Refluxing 30.6 g. 2,6-O2N(HO)C6H3Me in 150 cc. EtOH containing 4.6 g. Na 6 h. with 25.4 g. PhCH2Cl, adding H2O, distilling off the EtOH in vacuo, and extracting with Et2O give 63.8% 2,6-O2N(PhCH2O)C6H3Me (XIV), Bp. 8 170-6°, m. 65-6°. Condensation of XIV with (CO2Et)2 in the presence of EtOK gives the 2-nitro-6-benzoyloxyphenylpyruvic acid which is directly converted into 64% (overall) 4-benzoyl-2-indolecarboxylic acid (XV) (purified via its Na salt), m. 241-2°. Decarboxylation of XV in quinoline in the presence of Cu powder gives 62% 4-benzoyloxyindole (XVI), needles, m. 72-4°, which, treated in MeOH with H in the presence of IV, gives 4-hydroxyindole, hexagonal plates, m. 97-9°. Treating XVI with Me2NH in the same way as in the preparation of V gives 69% 4-benzoylgramine (XVII), hexagonal leaflets, m. 94-8°. Treating the methiodide of XVII with NaCl gives 60% 4-benzoyl-3-indoleacetoneitrile, m. 97-100°, which, reduced with LiAlH4, gives 81% 4-benzoyltryptamine, plates, m. 117-20° [acid oxalate (XVIII), hexagonal plates, m. 188-9°]. Shaking 3.3 g. XVIII in 270 cc. MeOH with 2% H and H gives 4-hydroxytryptamine (XIX) oxalate, clusters of platelets, m. 258-70°; free base does not crystallize. XIX-XIII complex, needles, m. 250-5°. Condensation of 121.5 g. 2,4-O2N(PhCH2O)C6H3Me with (CO2Et)2 gives 91% 2-nitro-6-benzoyloxyphenylpyruvic acid, m. 133-5° (B. and S. (loc. cit.) found 89-90°), which is converted into 51% 6-benzoyl-2-indolecarboxylic acid (XX), m. 199-200° (decomposition). Decarboxylation of XX gives 4% 6-benzoyloxyindole, leaflets, m. 118-20°, which, with Pd and H in MeOH, gives 6-hydroxyindole (XXI),

hexagonal leaflets, m. 124-6°. XXI is converted into 80% 6-benzoyloxygramine (XXII), long rods, m. 136-8°. Converting XXII into the methiodide and treating the latter with NaOH gives 75% 6-benzoyloxy-3-indoleacetonitrile, leaflets, m. 136-7°, which, reduced with LiAlH₄ in THF, gives 71% 6-benzoyloxytryptamine (XXIII), fine needles, m. 92-6° (oxalate, shiny leaflets, m. 260-5°). XXIII, debenzoylated with Pd and H, gives 6-hydroxytryptamine (XXIV) which does not crystallize. XXIII is converted into its sulfate and the latter (1.4 g.) is shaken in 500 cc. H₂O with 500 mg. IV and H, the filtrate concentrated to 100 cc., and 0.72 g. XIII added, giving 85% XXIV-XIII complex, fine needles, m. 212-15°. The UV and IR absorption maximum of some of the compds. are given.

RX(1) OF 3



NOTE: Classification: Isomerisation; Reductive coupling; Cyclisation; Heterocycle formation; # Conditions: Na2S2O4 NaOH; H2O

L20 ANSWER 8 OF 8 CASREACT COPYRIGHT 2009 ACS on STN

43:8352 Chemistry of bacteria. I. Synthesis of hydroxyindoles. Beer, R. J. S.; Clarke, Kenneth; Khosra, H. G.; Robertson, Alexander Journal of the Chemical Society 1905-9 (Unavailable) 1948. CODEN: JCSOAB. ISSN: 0368-1769.


AB Indoles hydroxylated in the C6H6 nucleus are needed to identify 2 isomeric products (C8H7ON) obtained in the degradation of violaceum (from Chromobacterium violaceum). 2,5-O2N(HO)C6H3CHO (5 g.), 6 g. acetic acid, 6 g. AcONa, and 15 mL Ac2O, heated 1.5 h. on the steam bath, the mixture kept 10 h., and diluted with 10 mL MeOH and 5 mL H2O, give 5-keto-2-methyl-4-(2-nitro-5-acetoxybenzylidene)-4,5-dihydrooxazole, yellow, m. 140°, and the hydroxybenzylidene analog, m. 173°; addition of 20 mL concentrated H2SO4 to 25 mL H2O containing 4 g. of the

mixed

salts and boiling 1 min. give 3.2 g. (2-nitro-5-hydroxyphenyl)pyruvic acid (I), m. 194° (decomposition); I also results on boiling the salts 4-5 h. with 1.5 N HCl; oxime, m. 181°. I (5 g.) in 34 mL NH4OH (d. 0.88) and 14 mL H2O, treated with 44 g. FeSO4 in 48 mL H2O and the mixture refluxed 1.5 h., gives 3 g. 5-hydroxy-2-indolecarboxylic acid (II), m. 246° (decomposition); decarboxylation of 0.5 g. II in 5 mL C3H5(OH)3 at 225-30° (25 min.) gives a poor yield of 5-hydroxyindole (picrate, orange-red, m. 167°); other conditions did not give better results. 2,4-O2N(PhCH2O)C6H3COO2N (oxime, m.

182° (decomposition)) yields 6-(benzyloxy)-2-indolecarboxylic acid, m. 200° (decomposition), debenzoylation of which (2 g. and 2 g. 10% Pd-C in 100 mL AcOH) gives 0.9 g. 6-hydroxy-2-indolecarboxylic acid (III), m. 236° (decomposition); 0.5 g. III in C3H5(OH)3, heated 20 min. at 220-30°, gives 0.2 g. 6-hydroxyindole. m. 125.5° (picrate, dark red, m. 154-6° (decomposition)). 2,3-O2N(HO)C6H3Me (5 g.), 3.8 mL PhC6H5, and excess K2CO3 in 100 mL boiling Me2CO (2.5 h.) give 6.5 g. 2-nitro-3-(benzyloxy)toluene, m. 36°; condensation of 24 g. with 15 g. CO2Et/2 with EtOH gives 15.5 g. 2-nitro-3-(benzyloxy)phenylpyruvic acid (IV), yellow, m. 125° (oxime, m. 190-1°). Reduction and cyclization of 3.8 g. IV give 2.2 g. 7-(benzyloxy)-2-indolecarboxylic acid (V), m. 164°; debenzoylation of 2 g. V gives 1 g. 7-hydroxy-2-indolecarboxylic acid, m. 252°; decarboxylation of 0.5 g. yields 20 mg. 7-hydroxyindole, m. 96°. 2,6-O2N(HO)C6H3CHO (1 g.) and 0.4 g. NaNO2 in EtOH at -10°, treated with 0.7 g. KOH in 10 mL EtOH, the mixture carefully acidified with cooled HCl, and the yellow oil (obtained by extraction with ether) gently warmed (15 min.) with 2 g. AcONa and 3 mL Ac2O, give 1.35 g. β,2-dinitro-6-acetoxystyrene (VI), light tan, m. 128-30°. Reaction of 0.5 g. VI, 2 g. Fe, 4 mL EtOH, and 4 mL AcOH, gently refluxed 10-12 min., gives 0.1-0.12 g. 4-acetoxyindole (VII), m. 100°; 0.1 g. VII in 3 mL MeOH saturated at 0° with NH3 and the solution kept 16 h. below 5°, gives 70-80% 4-hydroxyindole, m. 98° (picrate, red, becomes partly yellow at 150° and chars at 180°); it gives a dark blue color with alc. FeCl3 and a red color and strong green fluorescence with Ehrlich reagent in the cold. 2,5-O2N(HO)C6H3CHO yields β,2-dinitro-5-acetoxystyrene, straw, m. 118-19°; this yields 50-58% 5-acetoxyindole, m. 113-14°, which gives 5-hydroxyindole. 5,2-O2N(HO)C6H3CHO (1 g.) and EtNO2 give 0.7 g. β,2-dinitro-5-acetoxy-β-methylstyrene, pale brown, m. 88-9°; reduction with Fe and AcOH gives 5-acetoxy-2-methylindole, m. 128-30°, which yields 80% 5-hydroxy-2-methylindole, m. 133-4°, gives a brownish purple color with alc. FeCl3 and a very intense red color with Ehrlich reagent (picrate, red, m. 157-8° (decomposition)). 6,2-O2N(HO)C6H3CHO (1 g.), 0.8 g. acetic acid, 0.8 g. AcONa, and 3 mL Ac2O, heated 1 h. on the steam bath, give 1.3 g. 5-keto-4-(2-nitro-6-acetoxybenzylidene)-2-methyl-4,5-dihydrooxazole, yellow, m. 180°; hydrolysis of 1 g. with 1.5 N HCl (6 h. on the steam bath) gives 0.49 g. 5-nitro-3-hydroxyoxazolin, yellow, m. 206-7° (2,4-dinitrophenylhydrazine, orange, m. above 250°). 6-Nitro-2-(benzyloxy)toluene, m. 62°. 2,6-O2N(HO)C6H3CHO (1 g.), 0.5 mL AcOH, 0.5 g. AcONa, and 5 mL EtOH gradually deposit 0.85 g. 2-(2-nitro-6-hydroxyphenyl)-5-nitro-1,3,2H-benzoxazine, orange-yellow, m. 204-5°; alc. FeCl3 gives an orange color slowly changing to red; acetate, m. 164°. The stability of the various hydroxyindoles is discussed.


RX(2) OF 2



NOTE: Classification: Isomerisation; Reductive coupling; Cyclisation; Heterocycle formation; # Conditions: FeSO4 NH4OH; H2O; R14468

Print selected from 10562215.trn

=>
Uploading C:\Program Files\Stnexp\Queries\10562215-9999.str



chain nodes :
7 8 9
ring nodes :
1 2 3 4 5 6
ring/chain nodes :
10
chain bonds :
5-8 6-7 8-9 9-10
ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6
exact/norm bonds :
9-10
exact bonds :
5-8 6-7 8-9
normalized bonds :
1-2 1-6 2-3 3-4 4-5 5-6

Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS 9:CLASS 10:CLASS

L21 STRUCTURE UPLOADED

=> file caplus
FILE 'CAPLUS' ENTERED AT 11:33:50 ON 16 MAR 2009
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The ACS Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 16 Mar 2009 VOL 150 ISS 12

Print selected from 10562215.trn

FILE LAST UPDATED: 15 Mar 2009 (20090315/ED)

Caplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 121
REGISTRY INITIATED
Substance data SEARCH and crossover from CAS REGISTRY in progress...
Use DISPLAY HITSTR (or PHITSTR) to directly view retrieved structures.

SAMPLE SEARCH INITIATED 11:33:53 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 639 TO ITERATE

100.0% PROCESSED	639 ITERATIONS	50 ANSWERS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)		
SEARCH TIME: 00.00.01		

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 11264 TO 14296
PROJECTED ANSWERS: 7023 TO 9457

L22 50 SEA SSS SAM L21

L23 14 L22

=> file reg
FILE 'REGISTRY' ENTERED AT 11:33:58 ON 16 MAR 2009
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 13 MAR 2009 HIGHEST RN 1120564-02-4
DICTIONARY FILE UPDATES: 13 MAR 2009 HIGHEST RN 1120564-02-4

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 9, 2009.

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and

Print selected from 10562215.trn

predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stdoc/properties.html>

=> s l21
SAMPLE SEARCH INITIATED 11:34:00 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 639 TO ITERATE

100.0% PROCESSED 639 ITERATIONS 50 ANSWERS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 11264 TO 14296
PROJECTED ANSWERS: 7023 TO 9457

L24 50 SEA SSS SM L21

=> s l21 full
FULL SEARCH INITIATED 11:34:07 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 13107 TO ITERATE

100.0% PROCESSED 13107 ITERATIONS 8317 ANSWERS
SEARCH TIME: 00.00.01

L25 8317 SEA SSS FUL L21

=> d his

(FILE 'HOME' ENTERED AT 11:15:49 ON 16 MAR 2009)

FILE 'CASREACT' ENTERED AT 11:16:04 ON 16 MAR 2009

L1 STRUCTURE UPLOADED
L2 3 S L1
L3 114 S L1 FULL
L4 0 S L3 AND CARBON MONOXIDE
L5 5949 S CARBON MONOXIDE

FILE 'REGISTRY' ENTERED AT 11:17:25 ON 16 MAR 2009
L6 1 S CARBON MONOXIDE/CN

FILE 'CASREACT' ENTERED AT 11:17:52 ON 16 MAR 2009

L7 10978 S 630-08-3
L8 2 S L7 AND L3
L9 STRUCTURE UPLOADED
L10 6 S L9
L11 151 S L9 FULL
L12 3 S L11 AND L7
L13 1 S L12 NOT L8
L14 3 S L11 AND IRON
L15 3 S L11 AND CARBONYL
L16 2 S L11 AND COMPLEX
L17 9 S L13-L16
L18 8 S L14-L16
L19 8 S L18 NOT L12
L20 8 S L18 NOT L8

Print selected from 10562215.trn

L21 STRUCTURE UPLOADED

FILE 'CAPLUS' ENTERED AT 11:33:50 ON 16 MAR 2009
S L21

FILE 'REGISTRY' ENTERED AT 11:33:53 ON 16 MAR 2009
L22 50 S L21

FILE 'CAPLUS' ENTERED AT 11:33:54 ON 16 MAR 2009
L23 14 S L22

FILE 'REGISTRY' ENTERED AT 11:33:58 ON 16 MAR 2009
L24 50 S L21
L25 8317 S L21 FULL

=> file caplus
FILE 'CAPLUS' ENTERED AT 11:34:26 ON 16 MAR 2009
USE IS SUBJECT TO THE TERMS OF YOUR STM CUSTOMER AGREEMENT.
PLEASE SEE "HELP USABTERMS" FOR DETAILS.
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 16 Mar 2009 VOL 150 ISS 12
FILE LAST UPDATED: 15 Mar 2009 (20090315/ED)

Caplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> help roles

CAS roles are CAS indexing terms consisting of codes that describe the new or novel information reported about a substance or a class of compounds. Specific roles have 3-letter codes. Super roles have 4-letter codes. Super roles are automatically generated from the specific roles, and are supported for searching. The PREP (Preparation) role is available for documents from 1907 to the present. Other roles are available for all indexed documents from 1967 to the present.

To search a role for a specific substance, append the CAS Registry Number or a Registry File L-number answer set with a slash and the code for the role, e.g., 47-68-5/THU. To search more than one role,

Print selected from 10562215.trn

separate a list of roles by commas and no spaces, e.g.,
67-68-5/THU,ADV. Only one role may be appended to an L-number answer
set. Use the OR operator to apply multiple roles to an L-number,
e.g., S LI/THU OR LI/ADV.

To search roles assigned to index headings for classes of compounds,
follow the heading with a slash and the role or roles separated by
commas, e.g., PHENOLS/POL,REM.

Roles are displayed in the RL (Role) field within the IT (Index Term)
field. Roles are included in any display format that contains the IT
or RL field. Enter SET ROLES OFF at an arrow prompt (→) to suppress
display of codes and text for roles. Enter SET ROLES CODES to
display only codes. Enter SET ROLES TEXT to return to default
display (codes and names). Enter HELP SET ROLES at an arrow prompt
for more information.

Enter HELP THESAURUS and HELP PCODE at an arrow prompt in this file
for information on using the role thesaurus to find role definitions
and narrower and broader terms.

The following is a hierarchical list of CAS roles. Under each super
role are listed the specific roles that generate the super role.

List of CAS Roles (1)

ANST Analytical Study

ANT Analyte
ANX Analytical Matrix
ARG Analytical Reagent Use
ARU Analytical Role, Unclassified

BIOL Biological Study

ADV Adverse Effect, Including Toxicity
AGR Agricultural Use
BAC Biological Activity or Effector, Except Adverse (2)
BCP Biochemical Process (3)
BMF Biomedical Manufacture
BOC Biological Occurrence (2)
BPN Biosynthetic Preparation
BPR Biological Process (2)
BSU Biological Study, Unclassified
BTU Biological Use, Unclassified
COS Cosmetic Use (3)
DGN Diagnostic Use (3)
DMA Drug Mechanism of Action (3)
FFD Food or Feed Use
MFU Metabolic Formation (2)
NPO Natural Product Occurrence (3)
PAC Pharmacokinetics (3)
PKT Pharmacokinetics (3)
THU Therapeutic Use

CMST Combinatorial Study (3)

CPN Combinatorial Preparation (3)

Print selected from 10562215.trn

CRT Combinatorial Reactant (3)
CRG Combinatorial Reagent (3)
CST Combinatorial Study (3)
CUS Combinatorial Use (3)

FORM Formation, Nonpreparative

FNU Formation, Unclassified
GFM Geological or Astronomical Formation
MFU Metabolic Formation (2)

OCU Occurrence

BOC Biological Occurrence (2)
GOC Geological or Astronomical Occurrence
NPO Natural Product Occurrence (3)
OCU Occurrence, Unclassified
POL Pollutant

PREP Preparation (4)

BMF Biomedical Manufacture
BPN Biosynthetic Preparation
BYP Byproduct
CPN Combinatorial Preparation (3)
IMF Industrial Manufacture
PUR Purification or Recovery
PNU Preparation, Unclassified (5)
SPN Synthetic Preparation

PROC Process

BCP Biochemical Process (3)
BPR Biological Process (2)
GPR Geological or Astronomical Process
PEP Physical, Engineering, or Chemical Process
CPS Chemical Process (6)
EPR Engineering Process (6)
PTP Physical Process (6)
REM Removal or Disposal

PRPH Prophetic Substance (7)

RACI Reactant or Reagent (2,6)

RCT Reactant (8)
CRI Combinatorial Reactant (3)
RST Reagent (3)
CRG Combinatorial Reagent (3)

USES Uses

AGR Agricultural Use
ARG Analytical Reagent Use
BTU Biological Use, Unclassified
CAT Catalyst Use
COS Cosmetic Use (3)

Print selected from 10562215.trn

CUS Combinatorial Use (3)
DGN Diagnostic Use (3)
FFD Food or Feed Use
MDA Modifier or Additive Use
NUU Other Use, Unclassified (9)
POF Polymer in Formulation
TEM Technical or Engineered Material Use
THU Therapeutic Use

Specific roles that are not upposted to any super roles:

NSC Miscellaneous
PRP Properties

- (1) Super roles have 4-letter codes. Specific roles have 3-letter codes. Under each super role are listed the corresponding specific roles that are retrieved when you search that super role.
- (2) Used from CA Vol. 66 (1967) to Vol. 135 (2001)
- (3) Used starting with CA Vol. 136 (2002)
- (4) The PRPP super role has been added to records back to 1907.
- (5) Used from CA vol. 66 (1967) to vol. 145 (2006).
- (6) Used from CA vol. 136 (2002) to CA vol. 145 (2006).
- (7) Used starting with records from CA vol. 148 (2008).
- (8) Searching the RCT (Reactant) role retrieves references from CA Vol. 66 (1967) to the present. Searching the RACT (Reactant or Reagent) super role retrieves references with the CRT, CRS, RGT, or RCT references starting with CA Vol. 136 (2002).
- (9) Starting with CA Vol. 136 (2002), the searchable text for the NUU role changed from NONBIOLOGICAL USE, UNCLASSIFIED/RL to OTHER USE, UNCLASSIFIED/RL. Search the code NUU/RL to retrieve records from CA Vol. 66 (1967) to the present.

=> d his

(FILE 'HME' ENTERED AT 11:15:49 ON 16 MAR 2009)

FILE 'CASREACT' ENTERED AT 11:16:04 ON 16 MAR 2009

L1 STRUCTURE UPLOADED
L2 3 S L1
L3 114 S L1 FULL
L4 0 S L3 AND CARBON MONOXIDE
L5 5849 S CARBON MONOXIDE

L6 FILE 'REGISTRY' ENTERED AT 11:17:25 ON 16 MAR 2009
1 S CARBON MONOXIDE/CN

L7 FILE 'CASREACT' ENTERED AT 11:17:52 ON 16 MAR 2009
10878 S 610-08-0

L8 2 S L7 AND L3
L9 STRUCTURE UPLOADED
L10 6 S L9
L11 151 S L9 FULL
L12 3 S L11 AND L7
L13 1 S L12 NOT L8
L14 3 S L11 AND IRON
L15 3 S L11 AND CARBONYL
L16 2 S L11 AND COMPLEX

Print selected from 10562215.trn

L17 9 S L13-L16
L18 8 S L14-L16
L19 8 S L18 NOT L12
L20 8 S L18 NOT L8
L21 STRUCTURE UPLOADED

FILE 'CAPLUS' ENTERED AT 11:53:50 ON 16 MAR 2009
S L21

L22 FILE 'REGISTRY' ENTERED AT 11:33:53 ON 16 MAR 2009
50 S L21

L23 FILE 'CAPLUS' ENTERED AT 11:33:54 ON 16 MAR 2009
14 S L22

L24 FILE 'REGISTRY' ENTERED AT 11:33:58 ON 16 MAR 2009
50 S L21
L25 6517 S L21 FULL

FILE 'CAPLUS' ENTERED AT 11:34:26 ON 16 MAR 2009

=> s l25/ract
2918 L25
3247332 RACT/RL
L26 1497 L25/RACT
(L25 (L) RACT/RL)

=> s l26 and carbon monoxide
1444457 CARBON
30215 CARBONS
1455082 CARBON
(CARBON OR CARBONS)
201108 MONOXIDE
1090 MONOXIDES
201684 MONOXIDE
(MONOXIDE OR MONOXIDES)
170555 CARBON MONOXIDE
(CARBON (W) MONOXIDE)
L27 5 L26 AND CARBON MONOXIDE

=> d cdb abs hitstr 1-
YOU HAVE REQUESTED DATA FROM 5 ANSWERS - CONTINUE? Y/(N):y

L27 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2009 ACS ON STN
2008:994205 Document No. 149:3066210 Iodomethane. Sulikowski, Gary A.;
Sulikowski, Michelle K.; Haukaas, Michael H.; Ebon, Bonglin (USA). e-BROS
Encyclopedia of Reagents for Organic Synthesis, No pp. given. John Wiley
& Sons, Ltd.: Chichester, UK. (English) 2001. CODEN: 99K0U1. OTHER
SOURCES: CASREACT 149:306621.

AB A review of the article Iodomethane.
IT 4212-33-3
RL: RCT (Reactant); RACT (Reactant or reagent)
(Iodomethane)

RN 4212-33-3 CAPLUS
CF Ethanone, 2-(2-nitrophenyl)-1-phenyl- (CA INDEX NAME)



127 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2009 ACS on STN
2008:510694 Document No. 349:794360 Palladium-catalyzed synthesis of 3-indolecarboxylic acid derivatives. Soderberg, Bjorn C. G.; Banini, Serge R.; Turner, Michael R.; Winter, Aaron R.; Arrington, Amanda K. (C. Eugene Bennett Department of Chemistry, West Virginia University, Morgantown, WV, 26506-6045, USA). Synthesis (6), 903-912 (English) 2008. CODEN: SYNTAS. ISSN: 0039-7881. OTHER SOURCES: CASREACT 149:79436. Publisher: Georg Thieme Verlag.

GI



AB Indoles having an ester functionality in the 3-position, e.g., I, were prepared from 2-(2-nitrophenyl)propenoic acid derivs. via a palladium-catalyzed reductive N-heteroannulation using carbon monoxide as the ultimate reducing agent. The starting materials were prepared either by a Stille coupling of 2-halo-1-nitrobenzenes with Et 2-(tributylstannyl)-2-propenoate or by vicarious nucleophilic substitution of nitrobenzenes followed by a Knoevenagel-type condensation with an aldehyde. Synthesis of an example of a 3-nitrile- and a 3-sulfone-substituted indole is also described using the same type of methodologies.

II 30095-98-8 81327-28-8

RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of (aza)indole carboxylates/sulfone/nitrile via palladium/phenanthroline (or tri-Ph phosphine) catalyzed reductive N-heteroannulation of α,β -unsatd. ester/sulfone/nitrile intermediates)


RN 30095-98-8 CAPLUS

CN Benzenecarboxylic acid, 2-nitro-, methyl ester (CA INDEX NAME)



RN 81327-28-8 CAPLUS

CN Benzenecarboxylic acid, 5-chloro-2-nitro-, 1,1-dimethylethyl ester (CA INDEX NAME)



II 344449-11-2P 494836-60-1P 878672-60-7P

1033265-16-5P 1033265-29-0P 1033265-31-0P

1033265-33-0P 1033265-35-0P 1033265-37-0P

1033265-39-2P 1033265-41-6P 1033265-69-8P


RL: RCT (Reactant); SFN (Synthetic preparation); PREP (Preparation);

RACT (Reactant or reagent)

(preparation of (aza)indole carboxylates/sulfone/nitrile via palladium/phenanthroline (or tri-Ph phosphine) catalyzed reductive N-heteroannulation of α,β -unsatd. ester/sulfone/nitrile intermediates)


RN 344449-11-2 CAPLUS

CN 2-Naphthalenecarboxylic acid, 1-nitro-, 1,1-dimethylethyl ester (CA INDEX NAME)



RN 494836-60-1 CAPLUS


CN Benzenecarboxylic acid, o-methylene-2-nitro-, methyl ester (CA INDEX NAME)




RN 878672-60-7 CAPLUS

CN Benzenecarboxylic acid, 5-bromo-2-nitro-, 1,1-dimethylethyl ester (CA INDEX NAME)

Print selected from 10562215.trn




RN 1033265-16-5 CAPLUS
CN Benzenecarboxylic acid, 5-methoxy-2-nitro-, 1,1-dimethylethyl ester (CA INDEX NAME)



RN 1033265-29-0 CAPLUS
CN Benzenecarboxylic acid, 2-nitro- α -(phenylmethylene)-, methyl ester (CA INDEX NAME)




RN 1033265-31-4 CAPLUS
CN Benzenecarboxylic acid, 5-bromo- α -methylene-2-nitro-, 1,1-dimethylethyl ester (CA INDEX NAME)




RN 1033265-33-6 CAPLUS
CN Benzenecarboxylic acid, 5-chloro- α -methylene-2-nitro-, 1,1-dimethylethyl

Print selected from 10562215.trn


ester (CA INDEX NAME)




RN 1033265-35-8 CAPLUS
CN Benzenecarboxylic acid, 5-methoxy- α -methylene-2-nitro-, 1,1-dimethylethyl ester (CA INDEX NAME)



RN 1033265-37-0 CAPLUS
CN 2-Naphthaleneacetic acid, α -methylene-1-nitro-, 1,1-dimethylethyl ester (CA INDEX NAME)




RN 1033265-39-2 CAPLUS
CN Benzenecarboxylic acid, 4-methoxy- α -methylene-2-nitro-, ethyl ester (CA INDEX NAME)



RN 1033265-41-6 CAPLUS

Print selected from 10562215.trn


CN Benzeneacetic acid, 2-(methoxycarbonyl)- α -methylene-6-nitro-, ethyl ester (CA INDEX NAME)



RN 1033265-69-8 CAPLUS

CN Benzeneacetic acid, α -butylidene-2-nitro-, ethyl ester, (aZ)- (CA INDEX NAME)

Double bond geometry as shown.



L27 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2009 ACS on STN

2007:404312 Document No. 146:462149 Development of novel catalytic reactions of heterocyclic compounds toward new pesticide development. Tanaka, Norio (Electr. Mater. Res. Lab., Nissan Chemical Industries, Ltd., Japan). Farmashia, 43(3), 231-235 (Japanese) 2007. CODEN: FARUAW. ISSN: 0014-8601. Publisher: Pharmaceutical Society of Japan.

AB A review on catalytic reactions of N-containing heterocyclic compds. for the development of pesticides by Nissan Chemical Industries, Ltd., Japan: (1) preparation of pyrazolocarboxylic acids by liquid-phase oxidation using Co Mn bromide catalysts, (2) N-alkylation of azoles with alcs. using Ru catalysts, (3) preparation of indoles by reductive cyclization of (E)- β -nitrostyrenes or o-nitrophenylacetones using transition metal catalysts in CO atmospheric

IT 1969-72-8D, o-Nitrophenylacetone, derivs.

RU: RCT (Reactant); RACT (Reactant or reagent)

(preparation of indoles by reductive cyclization of nitrostyrenes or nitrophenylacetones using transition metal catalysts in CO atmospheric)

RN 1969-72-8 CAPLUS

CN 2-Propanone, 1-(2-nitrophenyl)- (CA INDEX NAME)




Print selected from 10562215.trn


L27 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2009 ACS on STN

2005:14373 Document No. 142:936770 Process for the preparation of indoles via reductive cyclization. Sakurai, Yasuhiro; Utsunomiya, Tomohisa; Tanaka, Norio (Nissan Chemical Industries, Ltd., Japan). PCT Int. Appl. WO 2005/00812 A1 2005/0106, 31 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BE, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, KE, KG, KN, KL, LC, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, ST, SV, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW; R0: AT, BE, BF, BJ, CF, CG, CH, CL, CM, CY, DE, DK, ES, FI, FR, GB, GE, GR, IE, IT, LU, MC, MD, ME, NL, PT, SE, SN, TD, TG, TR. (Japanese). CODEN: PIIKX2. APPLICATION: WO 2004-JP9001 20040623. PRIORITY: JP 2003-184359 20030627.

GI



I




II


AB A process for the preparation of indoles I [R1, R2 = H, (un)substituted alkyl, etc.; R3 = (un)substituted alkyl, etc.; n = 0-4] from 2-nitrobenzyl ketone compds. II [R1, R2 = H, (un)substituted alkyl, etc.; R3 = (un)substituted alkyl, etc.; n = 0-4] in the presence of a catalyst comprising a group VIII metal of the periodic table was provided. For example, in a stainless autoclave, a mixture of 4-fluoro-2-nitrophenylacetone (1.0 g), carbon monoxide (50 kgf/cm2) at 120 °C for 5 h, afforded 4-fluoro-2-methylindole (0.64 g, 95.0% yield), analysed by liquid chromatog. The process enables an indole compound to be selectively produced in high yield from a 2-nitrobenzylcarbonyl compound, and hardly yields an indoline compound as a reduction byproduct, this byproduct generation having been a problem in the catalytic hydrogenation method employing a noble-metal catalyst. The indole derivative produced by the process is useful as an intermediate for various fine chems. such as medicines and agricultural chems.

Print selected from 10562215.trn

IT 39616-99-4P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation);
RACT (Reactant or reagent)
(preparation of 1-(4-fluoro-2-nitrophenyl)acetone via decarboxylation)
RN 39616-99-4 CAPLUS
CN 2-Propanone, 1-(4-fluoro-2-nitrophenyl)- (CA INDEX NAME)



IT 816450-30-3P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation);
RACT (Reactant or reagent)
(preparation of 2-(2-nitrophenyl)-3-oxobutanoic acid Me ester derivative via
nucleophilic substitution of 2,5-difluoronitrobenzene with Et
acetooacetate)
RN 816450-30-3 CAPLUS
CN Benzeneacetic acid, α -acetyl-4-fluoro-2-nitro-, methyl ester (CA
INDEX NAME)




IT 1969-72-8 6127-15-7 57330-58-2
81327-26-6 85355-52-8 606092-02-8
816450-35-8
RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of indole derivative from 2-nitrobenzyl ketone compds. using
group VIII metal catalysts.)
RN 1969-72-8 CAPLUS
CN 2-Propanone, 1-(2-nitrophenyl)- (CA INDEX NAME)




RN 6127-15-7 CAPLUS
CN 2-Propanone, 1-(4-bromo-2-nitrophenyl)- (CA INDEX NAME)


Print selected from 10562215.trn



RN 57330-58-2 CAPLUS
CN 2-Propanone, 1-[2-nitro-4-(trifluoromethyl)phenyl]- (CA INDEX NAME)



RN 81327-26-6 CAPLUS
CN 2-Propanone, 1-(5-chloro-2-nitrophenyl)- (CA INDEX NAME)




RN 85355-52-8 CAPLUS
CN 2-Butanone, 3-(2-nitrophenyl)- (CA INDEX NAME)




RN 606092-02-8 CAPLUS
CN 3-Pentan-2-one, 3-(4-fluoro-2-nitrophenyl)-4-hydroxy- (CA INDEX NAME)


Print selected from 10562215.tzn



RN 816450-35-8 CAPLUS
CN 2-Propanone, 1-[(4-fluoro-2-nitro-5-(phenylmethoxy)phenyl)]- (CA INDEX NAME)



IT 606092-03-9P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation);
RACT (Reactant or reagent)
(preparation of indole derivative from 2-nitrobenzyl ketone compds. using
group VIII metal catalysts.)
RN 606092-03-9 CAPLUS
CN Benzeneacetic acid, 4-fluoro- α -(1-hydroxyethylidene)-2-nitro-,
methyl ester (CA INDEX NAME)



127 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2009 ACS on STN
1981:442811 Document No. 95:428110 Original Reference No. 95:7321a,7324a
Catalytic synthesis of 3-substituted indoles using carbon
monoxide as building block and supported rhodium as catalyst.
Ucciani, Eugene; Bonfand, Andre (Lab. Chim. Org. Appl., Fac. Sci. Tech.,
Marseille, 13397/13, Fr.). Journal of the Chemical Society, Chemical
Communications (3), 82-3 (English) 1981. CODEN: JOCCAT. ISSN: 0022-4936.
OTHER SOURCES: CASREACT 95:42811.
AB Skatole was prepared (approx.70%) from 2-(2NOC6H4CH=CH2, under
hydroformylation conditions, (CO, H, 160°) using C-supported Rh as
a catalyst, via 2-(2NOC6H4CMeCHO formation by homogeneous catalysis, reduction

Print selected from 10562215.tzn

of the NO2 group by heterogeneous catalysis, cyclization and thermal
dehydration.
IT 78266-67-8P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation);
RACT (Reactant or reagent)
(preparation and reduction of, rhodium-catalyzed)
RN 78266-67-8 CAPLUS
CN Benzeneacetaldehyde, α -methyl-2-nitro- (CA INDEX NAME)



=> s ucciani/au
L28 0 UCCIANTI/AU
=> s ucciani ?/au
L29 318 UCCIANTI ?/AU
=> s l19 and rhodium
8 L19
78733 RHODIUM
33 RHODIUMS
78735 RHODIUM
(RHODIUM OR RHODIUMS)
L30 1 L19 AND RHODIUM
=> s bonfand/au
L31 0 BONFAND/AU
=> s bonfand ?/au
L32 33 BONFAND ?/AU
=> s l32 and rhodium
78733 RHODIUM
33 RHODIUMS
78735 RHODIUM
(RHODIUM OR RHODIUMS)
L33 2 L32 AND RHODIUM

=> s l33 not l27
MISSING OPERATOR L33 MOD
The search profile that was entered contains terms or
nested terms that are not separated by a logical operator.
=> s l33 not l27
L34 1 L33 NOT L27
=> d chib abs hitatr 1-
YOU HAVE REQUESTED DATA FROM 1 ANSWERS - CONTINUE? Y/(N):y

Print selected from 10562215.trn

L34 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2009 ACS on STN
1991:494766 Document No. 115:94766 Original Reference No. 115:16301a,16304a
Hydroformylation of fatty acid vinyl esters. Bonfand, A.;
Ucciani, E.; Luciani, A.; Sambuc, E. (Lab. Chim. Org. Appl., CNRS, Fr.).
Revue Francaise des Corps Gras, 37(9-10), 281-7 (French) 1990. CODEN:
RFGGAE. ISSN: 0035-3000.
AB Hydroformylation of vinyl esters of lauric, myristic, palmitic, and
stearic acids to obtain fatty esters of propanediol was investigated. Of
several catalytic systems tested, Rh/C was the most efficient, leading to
aldesters instead of alc. esters. Catalytic hydroformylation by Rh/C gave
only one aldester, ROOCH(CH3)CHO. Reduction by NaBH4 yielded 2 alc. esters,
ROOCH(CH3)CH2OH (minor product) and ROOCH2CH2CH(CH3)OH (major product),
regardless of the vinyl ester used.

=> d his

(FILE 'HOME' ENTERED AT 11:15:49 ON 16 MAR 2009)
FILE 'CASREACT' ENTERED AT 11:16:04 ON 16 MAR 2009
L1 STRUCTURE UPLOADED
L2 3 S L1
L3 114 S L1 FULL
L4 0 S L3 AND CARBON MONOXIDE
L5 5949 S CARBON MONOXIDE

FILE 'REGISTRY' ENTERED AT 11:17:25 ON 16 MAR 2009
L6 1 S CARBON MONOXIDE/CN
FILE 'CASREACT' ENTERED AT 11:17:52 ON 16 MAR 2009
L7 10978 S 630-08-0
L8 2 S L7 AND L3
L9 STRUCTURE UPLOADED
L10 6 S L9
L11 151 S L9 FULL
L12 3 S L11 AND L7
L13 1 S L12 NOT L8
L14 3 S L11 AND IRON
L15 3 S L11 AND CARBONYL
L16 2 S L11 AND COMPLEX
L17 9 S L13-L16
L18 8 S L14-L16
L19 8 S L18 NOT L12
L20 8 S L18 NOT L8
L21 STRUCTURE UPLOADED

FILE 'CAPLUS' ENTERED AT 11:33:50 ON 16 MAR 2009
S L21
FILE 'REGISTRY' ENTERED AT 11:33:53 ON 16 MAR 2009
L22 50 S L21
FILE 'CAPLUS' ENTERED AT 11:33:54 ON 16 MAR 2009
L23 14 S L22
FILE 'REGISTRY' ENTERED AT 11:33:58 ON 16 MAR 2009
L24 50 S L21
L25 8317 S L21 FULL

Print selected from 10562215.trn

FILE 'CAPLUS' ENTERED AT 11:34:26 ON 16 MAR 2009
L26 1497 S L25/RAC
L27 5 S L26 AND CARBON MONOXIDE
L28 0 S UCCIANT/4U
L29 318 S UCCIANT 1/4U
L30 1 S L19 AND RHODIUM
L31 0 S BONFAND/4U
L32 33 S BONFAND 1/4U
L33 2 S L32 AND RHODIUM
L34 1 S L33 NOT L27
=> s 630-08-0/rn
159419 630-08-0
863 630-08-0D
L35 159661 630-08-0/RN
(630-08-0 (NOTL) 630-08-0D)

=> s l35 and l26
L36 4 L35 AND L26

=> d his
(FILE 'HOME' ENTERED AT 11:15:49 ON 16 MAR 2009)
FILE 'CASREACT' ENTERED AT 11:16:04 ON 16 MAR 2009
L1 STRUCTURE UPLOADED
L2 3 S L1
L3 114 S L1 FULL
L4 0 S L3 AND CARBON MONOXIDE
L5 5949 S CARBON MONOXIDE
FILE 'REGISTRY' ENTERED AT 11:17:25 ON 16 MAR 2009
L6 1 S CARBON MONOXIDE/CN
FILE 'CASREACT' ENTERED AT 11:17:52 ON 16 MAR 2009
L7 10978 S 630-08-0
L8 2 S L7 AND L3
L9 STRUCTURE UPLOADED
L10 6 S L9
L11 151 S L9 FULL
L12 3 S L11 AND L7
L13 1 S L12 NOT L8
L14 3 S L11 AND IRON
L15 3 S L11 AND CARBONYL
L16 2 S L11 AND COMPLEX
L17 9 S L13-L16
L18 8 S L14-L16
L19 8 S L18 NOT L12
L20 8 S L18 NOT L8
L21 STRUCTURE UPLOADED

FILE 'CAPLUS' ENTERED AT 11:33:50 ON 16 MAR 2009
S L21
FILE 'REGISTRY' ENTERED AT 11:33:53 ON 16 MAR 2009
L22 50 S L21
FILE 'CAPLUS' ENTERED AT 11:33:54 ON 16 MAR 2009
L23 14 S L22

Print selected from 10562215.trn

FILE 'REGISTRY' ENTERED AT 11:33:58 ON 16 MAR 2009
L24 50 S L21
L25 8317 S L21 FULL

FILE 'CAPLOS' ENTERED AT 11:34:26 ON 16 MAR 2009
L26 1497 S L25/RACT
L27 5 S L26 AND CARBON MONOXIDE
L28 0 S UCCULANI/AU
L29 318 S UCCULANI 2/AU
L30 1 S L19 AND RHODIUM
L31 0 S BONFAND/AU
L32 33 S BONFAND 2/AU
L33 2 S L32 AND RHODIUM
L34 1 S L33 NOT L27
L35 158661 S 630-08-0/RW
L36 4 S L35 AND L26

=> s 136 not 127